

Random Attractors for the Stochastic FitzHugh-Nagumo System on Unbounded Domains

Bixiang Wang *

Department of Mathematics, New Mexico Institute of Mining and Technology
Socorro, NM 87801, USA
Email: bwang@nmt.edu

Abstract

The existence of a random attractor for the stochastic FitzHugh-Nagumo system defined on an unbounded domain is established. The pullback asymptotic compactness of the stochastic system is proved by uniform estimates on solutions for large space and time variables. These estimates are obtained by a cut-off technique.

Key words. Stochastic FitzHugh-Nagumo system, random attractor, pullback attractor, asymptotic compactness.

MSC 2000. Primary 37L55. Secondary 60H15, 35B40, 35B41.

1 Introduction

In this paper, we investigate the asymptotic behavior of solutions of the following stochastic FitzHugh-Nagumo system defined on $\mathbb{R}^n$:

$$du + (\lambda u - \Delta u + \alpha v)dt = (f(x, u) + g)dt + \phi_1 dw_1, \quad x \in \mathbb{R}^n, \quad t > 0,$$

$$dv + (\delta v - \beta u)dt = hdt + \phi_2 dw_2, \quad x \in \mathbb{R}^n, \quad t > 0,$$

where λ, α, δ and β are positive constants, $g \in L^2(\mathbb{R}^n)$ and $h \in H^1(\mathbb{R}^n)$ are given, $\phi_1 \in H^2(\mathbb{R}^n) \cap W^{2,p}(\mathbb{R}^n)$ for some $p \geq 2$, $\phi_2 \in H^1(\mathbb{R}^n)$, f is a nonlinear function satisfying certain dissipative conditions, w_1 and w_2 are independent two-sided real-valued Wiener processes on a probability space which will be specified later.

*Supported in part by NSF grant DMS-0703521

It is known that the asymptotic behavior of a random system is determined by a pullback random attractor. The concept of random attractors was introduced in [12, 13], which is an analogue of global attractors for deterministic dynamical systems as studied in [5, 16, 23, 26, 29]. When PDEs are defined in *bounded* domains, the existence of random attractors has been investigated by many authors, see, e.g., [3, 10, 11, 12, 13] and the references therein. However, in the case of *unbounded* domains, the existence of random attractors is not well understood. It seems that the work [9] by Bates, Lu and Wang is the only existence result of random attractors for PDEs defined on unbounded domains, where a random attractor for the Reaction-Diffusion equation on $\mathbb{R}^n$ was established. In this paper, we will study such attractors for the stochastic FitzHugh-Nagumo system defined on unbounded domains.

Notice that Sobolev embeddings are not compact when domains are unbounded. This introduces a major obstacle for proving the existence of attractors for PDEs on unbounded domains. For some deterministic PDEs, such difficulty can be overcome by the energy equation approach that was developed by Ball in [6, 7] and used in [14, 15, 17, 20, 21, 25, 31]). Under certain circumstances, the tail-estimates method must be used to deal with the problem caused by the unboundedness of domains. This approach was developed in [30] for deterministic parabolic equations and used in [1, 2, 4, 19, 22, 24, 27, 28]. Recently, the authors of [9] extended the tail-estimates method to the stochastic parabolic equations. In this paper, we will use the idea of uniform estimates on the tails of solutions to investigate the asymptotic behavior of the stochastic FitzHugh-Nagumo system defined on $\mathbb{R}^n$.

This paper is organized as follows. In the next section, we review the pullback random attractors theory for random dynamical systems. In Section 3, we define a continuous random dynamical system for the stochastic FitzHugh-Nagumo system on $\mathbb{R}^n$. Then we derive the uniform estimates of solutions in Section 4, which include the uniform estimates on the tails of solutions. Finally, we establish the asymptotic compactness of the random dynamical system and prove the existence of a pullback random attractor.

In the sequel, we adopt the following notations. We denote by $\|\cdot\|$ and $(\cdot, \cdot)$ the norm and the inner product of $L^2(\mathbb{R}^n)$, respectively. The norm of a given Banach space X is written as $\|\cdot\|_X$. We also use $\|\cdot\|_p$ to denote the norm of $L^p(\mathbb{R}^n)$. The letters c and c_i ($i = 1, 2, \dots$) are generic

positive constants which may change their values from line to line or even in the same line.

2 Preliminaries

In this section, we recall some basic concepts related to random attractors for stochastic dynamical systems. The reader is referred to [3, 8, 11, 13] for more details.

Let $(X, \|\cdot\|_X)$ be a separable Hilbert space with Borel σ -algebra $\mathcal{B}(X)$, and $(\Omega, \mathcal{F}, P)$ be a probability space.

Definition 2.1. $(\Omega, \mathcal{F}, P, (\theta_t)_{t \in \mathbb{R}})$ is called a metric dynamical system if $\theta : \mathbb{R} \times \Omega \rightarrow \Omega$ is $(\mathcal{B}(\mathbb{R}) \times \mathcal{F}, \mathcal{F})$ -measurable, θ_0 is the identity on Ω , $\theta_{s+t} = \theta_t \circ \theta_s$ for all $s, t \in \mathbb{R}$ and $\theta_t P = P$ for all $t \in \mathbb{R}$.

Definition 2.2. A continuous random dynamical system (RDS) on X over a metric dynamical system $(\Omega, \mathcal{F}, P, (\theta_t)_{t \in \mathbb{R}})$ is a mapping

$$\Phi : \mathbb{R}^+ \times \Omega \times X \rightarrow X \quad (t, \omega, x) \mapsto \Phi(t, \omega, x),$$

which is $(\mathcal{B}(\mathbb{R}^+) \times \mathcal{F} \times \mathcal{B}(X), \mathcal{B}(X))$ -measurable and satisfies, for P -a.e. $\omega \in \Omega$,

- (i) $\Phi(0, \omega, \cdot)$ is the identity on X ;
- (ii) $\Phi(t + s, \omega, \cdot) = \Phi(t, \theta_s \omega, \cdot) \circ \Phi(s, \omega, \cdot)$ for all $t, s \in \mathbb{R}^+$;
- (iii) $\Phi(t, \omega, \cdot) : X \rightarrow X$ is continuous for all $t \in \mathbb{R}^+$.

Hereafter, we always assume that Φ is a continuous RDS on X over $(\Omega, \mathcal{F}, P, (\theta_t)_{t \in \mathbb{R}})$.

Definition 2.3. A random bounded set $\{B(\omega)\}_{\omega \in \Omega}$ of X is called tempered with respect to $(\theta_t)_{t \in \mathbb{R}}$ if for P -a.e. $\omega \in \Omega$,

$$\lim_{t \rightarrow \infty} e^{-\beta t} d(B(\theta_{-t} \omega)) = 0 \quad \text{for all } \beta > 0,$$

where $d(B) = \sup_{x \in B} \|x\|_X$.

Definition 2.4. Let $\mathcal{D}$ be a collection of random subsets of X . Then $\mathcal{D}$ is called inclusion-closed if $D = \{D(\omega)\}_{\omega \in \Omega} \in \mathcal{D}$ and $\tilde{D} = \{\tilde{D}(\omega) \subseteq X : \omega \in \Omega\}$ with $\tilde{D}(\omega) \subseteq D(\omega)$ for all $\omega \in \Omega$ imply that $\tilde{D} \in \mathcal{D}$.

Definition 2.5. Let $\mathcal{D}$ be a collection of random subsets of X and $\{K(\omega)\}_{\omega \in \Omega} \in \mathcal{D}$. Then $\{K(\omega)\}_{\omega \in \Omega}$ is called an absorbing set of Φ in $\mathcal{D}$ if for every $B \in \mathcal{D}$ and P -a.e. $\omega \in \Omega$, there exists $t_B(\omega) > 0$ such that

$$\Phi(t, \theta_{-t}\omega, B(\theta_{-t}\omega)) \subseteq K(\omega) \quad \text{for all } t \geq t_B(\omega).$$

Definition 2.6. Let $\mathcal{D}$ be a collection of random subsets of X . Then Φ is said to be $\mathcal{D}$ -pullback asymptotically compact in X if for P -a.e. $\omega \in \Omega$, $\{\Phi(t_n, \theta_{-t_n}\omega, x_n)\}_{n=1}^\infty$ has a convergent subsequence in X whenever $t_n \rightarrow \infty$, and $x_n \in B(\theta_{-t_n}\omega)$ with $\{B(\omega)\}_{\omega \in \Omega} \in \mathcal{D}$.

Definition 2.7. Let $\mathcal{D}$ be a collection of random subsets of X and $\{\mathcal{A}(\omega)\}_{\omega \in \Omega} \in \mathcal{D}$. Then $\{\mathcal{A}(\omega)\}_{\omega \in \Omega}$ is called a $\mathcal{D}$ -random attractor (or $\mathcal{D}$ -pullback attractor) for Φ if the following conditions are satisfied, for P -a.e. $\omega \in \Omega$,

- (i) $\mathcal{A}(\omega)$ is compact, and $\omega \mapsto d(x, \mathcal{A}(\omega))$ is measurable for every $x \in X$;
- (ii) $\{\mathcal{A}(\omega)\}_{\omega \in \Omega}$ is invariant, that is,

$$\Phi(t, \omega, \mathcal{A}(\omega)) = \mathcal{A}(\theta_t \omega), \quad \forall t \geq 0;$$

- (iii) $\{\mathcal{A}(\omega)\}_{\omega \in \Omega}$ attracts every set in $\mathcal{D}$, that is, for every $B = \{B(\omega)\}_{\omega \in \Omega} \in \mathcal{D}$,

$$\lim_{t \rightarrow \infty} d(\Phi(t, \theta_{-t}\omega, B(\theta_{-t}\omega)), \mathcal{A}(\omega)) = 0,$$

where d is the Hausdorff semi-metric given by $d(Y, Z) = \sup_{y \in Y} \inf_{z \in Z} \|y - z\|_X$ for any $Y \subseteq X$ and $Z \subseteq X$.

The following existence result on a random attractor for a continuous RDS can be found in [8, 13].

Proposition 2.8. *Let $\mathcal{D}$ be an inclusion-closed collection of random subsets of X and Φ a continuous RDS on X over $(\Omega, \mathcal{F}, P, (\theta_t)_{t \in \mathbb{R}})$. Suppose that $\{K(\omega)\}_{\omega \in \Omega}$ is a closed absorbing set of Φ in $\mathcal{D}$ and Φ is $\mathcal{D}$ -pullback asymptotically compact in X . Then Φ has a unique $\mathcal{D}$ -random attractor $\{\mathcal{A}(\omega)\}_{\omega \in \Omega}$ which is given by*

$$\mathcal{A}(\omega) = \bigcap_{\tau \geq 0} \overline{\bigcup_{t \geq \tau} \Phi(t, \theta_{-t}\omega, K(\theta_{-t}\omega))}.$$

In this paper, we will take $\mathcal{D}$ as the collection of all tempered random subsets of $L^2(\mathbb{R}^n) \times L^2(\mathbb{R}^n)$ and prove the stochastic FitzHugh-Nagumo system on $\mathbb{R}^n$ has a $\mathcal{D}$ -random attractor.

3 Stochastic FitzHugh-Nagumo system on $\mathbb{R}^n$

In this section, we discuss the existence of a continuous random dynamical system for the stochastic FitzHugh-Nagumo system defined on $\mathbb{R}^n$:

$$du + (\lambda u - \Delta u + \alpha v)dt = (f(x, u) + g)dt + \phi_1 dw_1, \quad x \in \mathbb{R}^n, \quad t > 0, \quad (3.1)$$

$$dv + (\delta v - \beta u)dt = hdt + \phi_2 dw_2, \quad x \in \mathbb{R}^n, \quad t > 0, \quad (3.2)$$

with the initial conditions:

$$u(x, 0) = u_0(x), \quad v(x, 0) = v_0(x), \quad x \in \mathbb{R}^n, \quad (3.3)$$

where λ, α, δ and β are positive constants, $g \in L^2(\mathbb{R}^n)$ and $h \in H^1(\mathbb{R}^n)$ are given, $\phi_1 \in H^2(\mathbb{R}^n) \cap W^{2,p}(\mathbb{R}^n)$ for some $p \geq 2$, $\phi_2 \in H^1(\mathbb{R}^n)$, w_1 and w_2 are independent two-sided real-valued Wiener processes on a probability space which will be specified below, and f is a nonlinear function satisfying the conditions, $\forall x \in \mathbb{R}^n$ and $s \in \mathbb{R}$,

$$f(x, s)s \leq -\alpha_1 |s|^p + \psi_1(x), \quad (3.4)$$

$$|f(x, s)| \leq \alpha_2 |s|^{p-1} + \psi_2(x), \quad (3.5)$$

$$\frac{\partial f}{\partial s}(x, s) \leq \beta, \quad (3.6)$$

$$\left| \frac{\partial f}{\partial x}(x, s) \right| \leq \psi_3(x), \quad (3.7)$$

where α_1, α_2 and β are positive constants, $\psi_1 \in L^1(\mathbb{R}^n) \cap L^\infty(\mathbb{R}^n)$, and $\psi_2 \in L^2(\mathbb{R}^n) \cap L^q(\mathbb{R}^n)$ with $\frac{1}{q} + \frac{1}{p} = 1$, and $\psi_3 \in L^2(\mathbb{R}^n)$.

In the sequel, we consider the probability space $(\Omega, \mathcal{F}, P)$ where

$$\Omega = \{\omega = (\omega_1, \omega_2) \in C(\mathbb{R}, \mathbb{R}^2) : \omega(0) = 0\},$$

$\mathcal{F}$ is the Borel σ -algebra induced by the compact-open topology of Ω , and P the corresponding Wiener measure on $(\Omega, \mathcal{F})$. Then we have

$$(w_1(t, \omega), w_2(t, \omega)) = \omega(t), \quad t \in \mathbb{R}.$$

Define the time shift by

$$\theta_t \omega(\cdot) = \omega(\cdot + t) - \omega(t), \quad \omega \in \Omega, \quad t \in \mathbb{R}.$$

Then $(\Omega, \mathcal{F}, P, (\theta_t)_{t \in \mathbb{R}})$ is a measurable dynamical system.

Consider the stationary solutions of the one-dimensional equations

$$dy_1 + \lambda y_1 dt = dw_1(t), \tag{3.8}$$

and

$$dy_2 + \delta y_2 dt = dw_2(t). \tag{3.9}$$

The solutions to (3.8) and (3.9) are given by

$$y_1(\theta_t \omega_1) = -\lambda \int_{-\infty}^0 e^{\lambda \tau} (\theta_t \omega_1)(\tau) d\tau, \quad t \in \mathbb{R},$$

and

$$y_2(\theta_t \omega_2) = -\delta \int_{-\infty}^0 e^{\delta \tau} (\theta_t \omega_2)(\tau) d\tau, \quad t \in \mathbb{R}.$$

Note that there exists a θ_t -invariant set $\tilde{\Omega} \subseteq \Omega$ of full P measure such that $y_j(\theta_t \omega_j)$ ($j = 1, 2$) is continuous in t for every $\omega \in \tilde{\Omega}$, and the random variable $|y_j(\omega_j)|$ is tempered (see, e.g., [8]). Therefore, it follows from Proposition 4.3.3 in [3] that there exists a tempered function $r(\omega) > 0$ such that

$$\sum_{j=1}^2 (|y_j(\omega_j)|^2 + |y_j(\omega_j)|^p) \leq r(\omega). \tag{3.10}$$

where $r(\omega)$ satisfies, for P -a.e. $\omega \in \Omega$,

$$r(\theta_t \omega) \leq e^{\frac{\eta}{2}|t|} r(\omega), \quad t \in \mathbb{R}, \tag{3.11}$$

where $\eta = \min\{\lambda, \delta\}$. Then it follows from (3.10)-(3.11) that, for P -a.e. $\omega \in \Omega$,

$$\sum_{j=1}^2 (|y_j(\theta_t \omega_j)|^2 + |y_j(\theta_t \omega_j)|^p) \leq e^{\frac{\eta}{2}|t|} r(\omega), \quad t \in \mathbb{R}. \tag{3.12}$$

Putting $z_j(\theta_t \omega) = \phi_j y_j(\theta_t \omega_j)$, by (3.8) and (3.9) we have

$$dz_1 + \lambda z_1 dt = \phi_1 dw_1,$$

and

$$dz_2 + \delta z_2 dt = \phi_2 dw_2.$$

Let $\tilde{u} = u(t) - z_1(\theta_t \omega)$ and $\tilde{v} = v(t) - z_2(\theta_t \omega)$, where (u, v) satisfies (3.1)-(3.3). Then for $(\tilde{u}, \tilde{v})$, we have

$$\frac{d\tilde{u}}{dt} + \lambda\tilde{u} - \Delta\tilde{u} + \alpha\tilde{v} = f(x, \tilde{u} + z_1(\theta_t \omega)) + g + \Delta z_1(\theta_t \omega) - \alpha z_2(\theta_t \omega), \quad (3.13)$$

and

$$\frac{d\tilde{v}}{dt} + \delta\tilde{v} - \beta\tilde{u} = h + \beta z_1(\theta_t \omega). \quad (3.14)$$

By a Galerkin method as in [18], it can be proved that if f satisfies (3.4)-(3.7), then for P -a.e. $\omega \in \Omega$ and for all $(\tilde{u}_0, \tilde{v}_0) \in L^2(\mathbb{R}^n) \times L^2(\mathbb{R}^n)$, system (3.13)-(3.14) has a unique solution $(\tilde{u}(\cdot, \omega, \tilde{u}_0), \tilde{v}(\cdot, \omega, \tilde{v}_0)) \in C([0, \infty), L^2(\mathbb{R}^n) \times L^2(\mathbb{R}^n))$ with $\tilde{u}(0, \omega, \tilde{u}_0) = \tilde{u}_0$ and $\tilde{v}(0, \omega, \tilde{v}_0) = \tilde{v}_0$. Further, the solution $(\tilde{u}(t, \omega, \tilde{u}_0), \tilde{v}(t, \omega, \tilde{v}_0))$ is continuous with respect to $(\tilde{u}_0, \tilde{v}_0)$ in $L^2(\mathbb{R}^n) \times L^2(\mathbb{R}^n)$ for all $t \geq 0$. Throughout this paper, we always write

$$u(t, \omega, u_0) = \tilde{u}(t, \omega, u_0 - z_1(\omega)) + z_1(\theta_t \omega), \quad v(t, \omega, v_0) = \tilde{v}(t, \omega, v_0 - z_2(\omega)) + z_2(\theta_t \omega). \quad (3.15)$$

Then (u, v) is a solution of problem (3.1)-(3.3) in some sense. We now define a mapping $\Phi : \mathbb{R}^+ \times \Omega \times (L^2(\mathbb{R}^n) \times L^2(\mathbb{R}^n)) \rightarrow L^2(\mathbb{R}^n) \times L^2(\mathbb{R}^n)$ by

$$\Phi(t, \omega, (u_0, v_0)) = (u(t, \omega, u_0), v(t, \omega, v_0)), \quad \forall (t, \omega, (u_0, v_0)) \in \mathbb{R}^+ \times \Omega \times (L^2(\mathbb{R}^n) \times L^2(\mathbb{R}^n)). \quad (3.16)$$

Note that Φ satisfies conditions (i), (ii) and (iii) in Definition 2.2. Therefore, Φ is a continuous random dynamical system associated with the stochastic FitzHugh-Nagumo system on $\mathbb{R}^n$. In the next two sections, we establish uniform estimates for Φ and prove the existence of a random attractor.

4 Uniform estimates of solutions

In this section, we derive uniform estimates on the solutions of the stochastic FitzHugh-Nagumo system defined on $\mathbb{R}^n$ when $t \rightarrow \infty$. These estimates are necessary for proving the existence of bounded absorbing sets and the asymptotic compactness of the random dynamical system associated with the system. Particularly, we will show that the tails of the solutions for large space variables are uniformly small when time is sufficiently large.

From now on, we always assume that $\mathcal{D}$ is the collection of all tempered subsets of $L^2(\mathbb{R}^n) \times L^2(\mathbb{R}^n)$ with respect to $(\Omega, \mathcal{F}, P, (\theta_t)_{t \in \mathbb{R}})$. The next lemma shows that Φ has an absorbing set in $\mathcal{D}$.

Lemma 4.1. *Assume that $g, h \in L^2(\mathbb{R}^n)$ and (3.4)-(3.7) hold. Then there exists $\{K(\omega)\}_{\omega \in \Omega} \in \mathcal{D}$ such that $\{K(\omega)\}_{\omega \in \Omega}$ is a closed absorbing set for Φ in $\mathcal{D}$, that is, for any $B = \{B(\omega)\}_{\omega \in \Omega} \in \mathcal{D}$ and P -a.e. $\omega \in \Omega$, there is $T_B(\omega) > 0$ such that*

$$\Phi(t, \theta_{-t}\omega, B(\theta_{-t}\omega)) \subseteq K(\omega) \quad \text{for all } t \geq T_B(\omega).$$

Proof. Taking the inner product of (3.13) with $\beta \tilde{u}$ in $L^2(\mathbb{R}^n)$ we find that

$$\begin{aligned} & \frac{1}{2} \beta \frac{d}{dt} \|\tilde{u}\|^2 + \lambda \beta \|\tilde{u}\|^2 + \beta \|\nabla \tilde{u}\|^2 + \alpha \beta(\tilde{v}, \tilde{u}) \\ &= \beta(f(x, \tilde{u} + z_1(\theta_t \omega)), \tilde{u}) + \beta(g, \tilde{u}) + \beta(\Delta z_1(\theta_t \omega), \tilde{u}) - \alpha \beta(z_2(\theta_t \omega), \tilde{u}). \end{aligned} \quad (4.1)$$

Taking the inner product of (3.14) with $\alpha \tilde{v}$ in $L^2(\mathbb{R}^n)$ we find that

$$\frac{1}{2} \alpha \frac{d}{dt} \|\tilde{v}\|^2 + \alpha \delta \|\tilde{v}\|^2 - \alpha \beta(\tilde{v}, \tilde{u}) = \alpha(h, \tilde{v}) + \alpha \beta(z_1(\theta_t \omega), \tilde{v}). \quad (4.2)$$

Adding (4.1) and (4.2), we obtain that

$$\begin{aligned} & \frac{1}{2} \frac{d}{dt} (\beta \|\tilde{u}\|^2 + \alpha \|\tilde{v}\|^2) + \lambda \beta \|\tilde{u}\|^2 + \alpha \delta \|\tilde{v}\|^2 + \beta \|\nabla \tilde{u}\|^2 \\ &= \beta(f(x, \tilde{u} + z_1(\theta_t \omega)), \tilde{u}) + \beta(g, \tilde{u}) + \beta(\Delta z_1(\theta_t \omega), \tilde{u}) \\ & \quad - \alpha \beta(z_2(\theta_t \omega), \tilde{u}) + \alpha(h, \tilde{v}) + \alpha \beta(z_1(\theta_t \omega), \tilde{v}). \end{aligned} \quad (4.3)$$

We now estimate every term on the right-hand side of (4.3). For the nonlinear term, by (3.4)-(3.5) we obtain that

$$\begin{aligned} (f(x, \tilde{u} + z_1(\theta_t \omega)), \tilde{u}) &= (f(x, \tilde{u} + z_1(\theta_t \omega)), \tilde{u} + z_1(\theta_t \omega)) - (f(x, \tilde{u} + z_1(\theta_t \omega)), z_1(\theta_t \omega)) \\ &\leq -\alpha_1 \int_{\mathbb{R}^n} |u|^p dx + \int_{\mathbb{R}^n} \psi_1(x) dx - \int_{\mathbb{R}^n} f(x, u) z_1(\theta_t \omega) dx \\ &\leq -\alpha_1 \int_{\mathbb{R}^n} |u|^p dx + \int_{\mathbb{R}^n} \psi_1(x) dx + \alpha_2 \int_{\mathbb{R}^n} |u|^{p-1} |z_1(\theta_t \omega)| dx + \int_{\mathbb{R}^n} |\psi_2| |z_1(\theta_t \omega)| dx \\ &\leq -\alpha_1 \|u\|_p^p + \|\psi_1\|_1 + \frac{1}{2} \alpha_1 \|u\|_p^p + c_1 \|z_1(\theta_t \omega)\|_p^p + \frac{1}{2} \|\psi_2\|^2 + \frac{1}{2} \|z_1(\theta_t \omega)\|^2 \\ &\leq -\frac{1}{2} \alpha_1 \|u\|_p^p + c_2 (\|z_1(\theta_t \omega)\|_p^p + \|z_1(\theta_t \omega)\|^2) + c_3. \end{aligned} \quad (4.4)$$

The second term on the right-hand side of (4.3) is bounded by

$$|\beta|(g, \tilde{u})| \leq \beta \|g\| \|\tilde{u}\| \leq \frac{1}{4} \lambda \beta \|\tilde{u}\|^2 + \frac{\beta}{\lambda} \|g\|^2. \quad (4.5)$$

For the third term on the right-hand side of (4.3), we have

$$|\beta|(\nabla z_1(\theta_t \omega), \nabla \tilde{u})| \leq \frac{1}{2} \beta \|\nabla z_1(\theta_t \omega)\|^2 + \frac{1}{2} \beta \|\nabla \tilde{u}\|^2 \quad (4.6)$$

Similarly, the remaining terms on the right-hand side of (4.3) is bounded by

$$\begin{aligned} & \alpha \beta |(z_2(\theta_t \omega), \tilde{u})| + \alpha |(h, \tilde{v})| + \alpha \beta |(z_1(\theta_t \omega), \tilde{v})| \\ & \leq \frac{1}{4} \lambda \beta \|\tilde{u}\|^2 + \frac{1}{\lambda} \alpha^2 \beta \|z_2(\theta_t \omega)\|^2 + \frac{1}{4} \alpha \delta \|\tilde{v}\|^2 + \frac{\alpha}{\delta} \|h\|^2 + \frac{1}{4} \alpha \delta \|\tilde{v}\|^2 + \frac{\alpha \beta^2}{\delta} \|z_1(\theta_t \omega)\|^2 \\ & \leq \frac{1}{4} \lambda \beta \|\tilde{u}\|^2 + \frac{1}{\lambda} \alpha^2 \beta \|z_2(\theta_t \omega)\|^2 + \frac{1}{2} \alpha \delta \|\tilde{v}\|^2 + \frac{\alpha}{\delta} \|h\|^2 + \frac{\alpha \beta^2}{\delta} \|z_1(\theta_t \omega)\|^2. \end{aligned} \quad (4.7)$$

Then it follows from (4.3)-(4.7) that

$$\begin{aligned} & \frac{d}{dt} (\beta \|\tilde{u}\|^2 + \alpha \|\tilde{v}\|^2) + \lambda \beta \|\tilde{u}\|^2 + \alpha \delta \|\tilde{v}\|^2 + \beta \|\nabla \tilde{u}\|^2 + \alpha_1 \|u\|_p^p \\ & \leq c_4 (\|z_1(\theta_t \omega)\|_p^p + \|z_1(\theta_t \omega)\|^2 + \|\nabla z_1(\theta_t \omega)\|^2 + \|z_2(\theta_t \omega)\|^2) + c_5. \end{aligned} \quad (4.8)$$

Note that $z_1(\theta_t \omega) = \phi_1 y_1(\theta_t \omega_1)$, $z_2(\theta_t \omega) = \phi_2 y_2(\theta_t \omega_2)$, $\phi_2 \in H^1(\mathbb{R}^n)$ and $\phi_1 \in H^2(\mathbb{R}^n) \cap W^{2,p}(\mathbb{R}^n)$.

Therefore, the right-hand of (4.8) is bounded by

$$c_6 \sum_{j=1}^2 (|y_j(\theta_t \omega_j)|^p + |y_j(\theta_t \omega_j)|^2) + c_7 = p_1(\theta_t \omega) + c_7. \quad (4.9)$$

By (3.12), we find that for P -a.e. $\omega \in \Omega$,

$$p_1(\theta_\tau \omega) \leq c_6 e^{\frac{1}{2} \eta |\tau|} r(\omega), \quad \forall \tau \in \mathbb{R}. \quad (4.10)$$

It follows from (4.8)-(4.9) that, for all $t \geq 0$,

$$\frac{d}{dt} (\beta \|\tilde{u}\|^2 + \alpha \|\tilde{v}\|^2) + \lambda \beta \|\tilde{u}\|^2 + \alpha \delta \|\tilde{v}\|^2 + \beta \|\nabla \tilde{u}\|^2 + \alpha_1 \|u\|_p^p \leq p_1(\theta_t \omega) + c_7, \quad (4.11)$$

which implies that, for all $t \geq 0$,

$$\frac{d}{dt} (\beta \|\tilde{u}\|^2 + \alpha \|\tilde{v}\|^2) + \lambda \beta \|\tilde{u}\|^2 + \alpha \delta \|\tilde{v}\|^2 \leq p_1(\theta_t \omega) + c_7.$$

Note that $\eta = \min\{\lambda, \delta\}$. We get that, for all $t \geq 0$,

$$\frac{d}{dt} (\beta \|\tilde{u}\|^2 + \alpha \|\tilde{v}\|^2) + \eta (\beta \|\tilde{u}\|^2 + \alpha \|\tilde{v}\|^2) \leq p_1(\theta_t \omega) + c_7. \quad (4.12)$$

By Gronwall's lemma, we find that, for all $t \geq 0$,

$$\begin{aligned} & \beta \|\tilde{u}(t, \omega, \tilde{u}_0(\omega))\|^2 + \alpha \|\tilde{v}(t, \omega, \tilde{v}_0(\omega))\|^2 \\ & \leq e^{-\eta t} (\beta \|\tilde{u}_0(\omega)\|^2 + \alpha \|\tilde{v}_0(\omega)\|^2) + \int_0^t e^{\eta(\tau-t)} p_1(\theta_\tau \omega) d\tau + \frac{c_7}{\eta}. \end{aligned} \quad (4.13)$$

By replacing ω by $\theta_{-t}\omega$, we get from (4.13) and (4.10) that, for all $t \geq 0$,

$$\begin{aligned} & \beta \|\tilde{u}(t, \theta_{-t}\omega, \tilde{u}_0(\theta_{-t}\omega))\|^2 + \alpha \|\tilde{v}(t, \theta_{-t}\omega, \tilde{v}_0(\theta_{-t}\omega))\|^2 \\ & \leq e^{-\eta t} (\beta \|\tilde{u}_0(\theta_{-t}\omega)\|^2 + \alpha \|\tilde{v}_0(\theta_{-t}\omega)\|^2) + \int_0^t e^{\eta(\tau-t)} p_1(\theta_{\tau-t}\omega) d\tau + \frac{c_7}{\eta}. \\ & \leq e^{-\eta t} (\beta \|\tilde{u}_0(\theta_{-t}\omega)\|^2 + \alpha \|\tilde{v}_0(\theta_{-t}\omega)\|^2) + \int_{-t}^0 e^{\eta\tau} p_1(\theta_\tau \omega) d\tau + \frac{c_7}{\eta}. \\ & \leq e^{-\eta t} (\beta \|\tilde{u}_0(\theta_{-t}\omega)\|^2 + \alpha \|\tilde{v}_0(\theta_{-t}\omega)\|^2) + c_6 \int_{-t}^0 e^{\frac{1}{2}\eta\tau} r(\omega) d\tau + \frac{c_7}{\eta}. \\ & \leq e^{-\eta t} (\beta \|\tilde{u}_0(\theta_{-t}\omega)\|^2 + \alpha \|\tilde{v}_0(\theta_{-t}\omega)\|^2) + \frac{2c_6}{\eta} r(\omega) + \frac{c_7}{\eta}. \end{aligned} \quad (4.14)$$

By (3.15) and (3.16) we have

$$\Phi(t, \omega, (u_0, v_0)(\omega)) = (\tilde{u}(t, \omega, u_0(\omega) - z_1(\omega)) + z_1(\theta_t \omega), \tilde{v}(t, \omega, v_0(\omega) - z_2(\omega)) + z_2(\theta_t \omega)).$$

So by (4.14) we get that, for all $t \geq 0$,

$$\begin{aligned} & \|\Phi(t, \theta_{-t}\omega, (u_0, v_0)(\theta_{-t}\omega))\|^2 \\ & = \|\tilde{u}(t, \theta_{-t}\omega, u_0(\theta_{-t}\omega) - z_1(\theta_{-t}\omega)) + z_1(\omega)\|^2 \\ & \quad + \|\tilde{v}(t, \theta_{-t}\omega, v_0(\theta_{-t}\omega) - z_2(\theta_{-t}\omega)) + z_2(\omega)\|^2 \\ & \leq 2\|\tilde{u}(t, \theta_{-t}\omega, u_0(\theta_{-t}\omega) - z_1(\theta_{-t}\omega))\|^2 + 2\|z_1(\omega)\|^2 \\ & \quad + 2\|\tilde{v}(t, \theta_{-t}\omega, v_0(\theta_{-t}\omega) - z_2(\theta_{-t}\omega))\|^2 + 2\|z_2(\omega)\|^2 \\ & \leq 2\left(\frac{1}{\alpha} + \frac{1}{\beta}\right) e^{-\eta t} (\beta \|u_0(\theta_{-t}\omega) - z_1(\theta_{-t}\omega)\|^2 + \alpha \|v_0(\theta_{-t}\omega) - z_2(\theta_{-t}\omega)\|^2) \end{aligned}$$

$$\begin{aligned}
& +2\|z_1(\omega)\|^2 + 2\|z_2(\omega)\|^2 + c_8 r(\omega) + c_9 \\
& \leq c_{10} e^{-\eta t} (\|u_0(\theta_{-t}\omega)\|^2 + \|v_0(\theta_{-t}\omega)\|^2 + \|z_1(\theta_{-t}\omega)\|^2 + \|z_2(\theta_{-t}\omega)\|^2) \\
& \quad + 2\|z_1(\omega)\|^2 + 2\|z_2(\omega)\|^2 + c_8 r(\omega) + c_9
\end{aligned} \tag{4.15}$$

Note that $\|z_1(\omega)\|^2$, $\|z_2(\omega)\|^2$ and $\{B(\omega)\}_{\omega \in \Omega} \in \mathcal{D}$ are tempered. Therefore, for $(u_0, v_0)(\theta_{-t}\omega) \in B(\theta_{-t}\omega)$, then there is $T_B(\omega) > 0$ such that for all $t \geq T_B(\omega)$,

$$c_{10} e^{-\eta t} (\|u_0(\theta_{-t}\omega)\|^2 + \|v_0(\theta_{-t}\omega)\|^2 + \|z_1(\theta_{-t}\omega)\|^2 + \|z_2(\theta_{-t}\omega)\|^2) \leq c_8 r(\omega) + c_9,$$

which along with (4.15) shows that, for all $t \geq T_B(\omega)$,

$$\|\Phi(t, \theta_{-t}\omega, (u_0, v_0)(\theta_{-t}\omega))\|^2 \leq 2 (\|z_1(\omega)\|^2 + \|z_2(\omega)\|^2 + c_8 r(\omega) + c_9). \tag{4.16}$$

Given $\omega \in \Omega$, denote by

$$K(\omega) = \{(u, v) \in L^2(\mathbb{R}^n) \times L^2(\mathbb{R}^n) : \|u\|^2 + \|v\|^2 \leq 2 (\|z_1(\omega)\|^2 + \|z_2(\omega)\|^2 + c_8 r(\omega) + c_9)\}.$$

Then $\{K(\omega)\}_{\omega \in \Omega} \in \mathcal{D}$. Further, (4.16) shows that $\{K(\omega)\}_{\omega \in \Omega}$ is an absorbing set for Φ in $\mathcal{D}$, which completes the proof. $\square$

Lemma 4.2. *Assume that $g, h \in L^2(\mathbb{R}^n)$ and (3.4)-(3.7) hold. Let $B = \{B(\omega)\}_{\omega \in \Omega} \in \mathcal{D}$. Then for every $T_1 \geq 0$ and P -a.e. $\omega \in \Omega$, $u(t, \omega, u_0(\omega))$ and $\tilde{u}(t, \omega, \tilde{u}_0(\omega))$ satisfy, for all $t \geq T_1$,*

$$\int_{T_1}^t e^{\eta(s-t)} \|u(s, \theta_{-t}\omega, u_0(\theta_{-t}\omega))\|_p^p ds \leq c e^{-\eta t} (\|\tilde{u}_0(\theta_{-t}\omega)\|^2 + \|\tilde{v}_0(\theta_{-t}\omega)\|^2) + c(1 + r(\omega)), \tag{4.17}$$

$$\int_{T_1}^t e^{\eta(s-t)} \|\nabla \tilde{u}(s, \theta_{-t}\omega, \tilde{u}_0(\theta_{-t}\omega))\|^2 ds \leq c e^{-\eta t} (\|\tilde{u}_0(\theta_{-t}\omega)\|^2 + \|\tilde{v}_0(\theta_{-t}\omega)\|^2) + c(1 + r(\omega)), \tag{4.18}$$

where c is a positive deterministic constant independent of T_1 , and $r(\omega)$ is the tempered function in (3.10).

Proof. First replacing t by T_1 and then replacing ω by $\theta_{-t}\omega$ in (4.13), we find that

$$\begin{aligned}
& \beta \|\tilde{u}(T_1, \theta_{-t}\omega, \tilde{u}_0(\theta_{-t}\omega))\|^2 + \alpha \|\tilde{v}(T_1, \theta_{-t}\omega, \tilde{v}_0(\theta_{-t}\omega))\|^2 \\
& \leq e^{-\eta T_1} (\beta \|\tilde{u}_0(\theta_{-t}\omega)\|^2 + \alpha \|\tilde{v}_0(\theta_{-t}\omega)\|^2) + \int_0^{T_1} e^{\eta(\tau-T_1)} p_1(\theta_{\tau-t}\omega) d\tau + c
\end{aligned}$$

Multiply the above by $e^{\eta(T_1-t)}$ and then simplify to get that

$$\begin{aligned} & e^{\eta(T_1-t)} (\beta \|\tilde{u}(T_1, \theta_{-t}\omega, \tilde{u}_0(\theta_{-t}\omega))\|^2 + \alpha \|\tilde{v}(T_1, \theta_{-t}\omega, \tilde{v}_0(\theta_{-t}\omega))\|^2) \\ & \leq e^{-\eta t} (\beta \|\tilde{u}_0(\theta_{-t}\omega)\|^2 + \alpha \|\tilde{v}_0(\theta_{-t}\omega)\|^2) + \int_0^{T_1} e^{\eta(\tau-t)} p_1(\theta_{\tau-t}\omega) d\tau + ce^{\eta(T_1-t)}. \end{aligned} \quad (4.19)$$

By (4.10), the second term on the right-hand side of (4.19) satisfies

$$\begin{aligned} & \int_0^{T_1} e^{\eta(\tau-t)} p_1(\theta_{\tau-t}\omega) d\tau = \int_{-t}^{T_1-t} e^{\eta\tau} p_1(\theta_\tau\omega) d\tau \\ & \leq c_6 r(\omega) \int_{-t}^{T_1-t} e^{\frac{1}{2}\eta\tau} d\tau \leq \frac{2}{\eta} c_6 r(\omega) e^{\frac{1}{2}\eta(T_1-t)}. \end{aligned} \quad (4.20)$$

From (4.19)-(4.20) it follows that

$$\begin{aligned} & e^{\eta(T_1-t)} (\beta \|\tilde{u}(T_1, \theta_{-t}\omega, \tilde{u}_0(\theta_{-t}\omega))\|^2 + \alpha \|\tilde{v}(T_1, \theta_{-t}\omega, \tilde{v}_0(\theta_{-t}\omega))\|^2) \\ & \leq e^{-\eta t} (\beta \|\tilde{u}_0(\theta_{-t}\omega)\|^2 + \alpha \|\tilde{v}_0(\theta_{-t}\omega)\|^2) + \frac{2}{\eta} c_6 r(\omega) e^{\frac{1}{2}\eta(T_1-t)} + ce^{\eta(T_1-t)}. \end{aligned} \quad (4.21)$$

Note that (4.11) implies, for all $t \geq 0$,

$$\frac{d}{dt} (\beta \|\tilde{u}\|^2 + \alpha \|\tilde{v}\|^2) + \eta (\beta \|\tilde{u}\|^2 + \alpha \|\tilde{v}\|^2) + \beta \|\nabla \tilde{u}\|^2 + \alpha_1 \|u\|_p^p \leq p_1(\theta_t\omega) + c.$$

Multiplying the above by $e^{\eta t}$ and then integrating over (T_1, t) , we get that, for all $t \geq T_1$,

$$\begin{aligned} & \beta \|\tilde{u}(t, \omega, \tilde{u}_0(\omega))\|^2 + \alpha \|\tilde{v}(t, \omega, \tilde{v}_0(\omega))\|^2 \\ & + \int_{T_1}^t e^{\eta(s-t)} \|\nabla \tilde{u}(s, \omega, \tilde{u}_0(\omega))\|^2 ds + \alpha_1 \int_{T_1}^t e^{\eta(s-t)} \|u(s, \omega, u_0(\omega))\|_p^p ds \\ & \leq e^{\eta(T_1-t)} (\beta \|\tilde{u}(T_1, \omega, \tilde{u}_0(\omega))\|^2 + \alpha \|\tilde{v}(T_1, \omega, \tilde{v}_0(\omega))\|^2) \\ & + \int_{T_1}^t e^{\eta(s-t)} p_1(\theta_s\omega) ds + c \int_{T_1}^t e^{\eta(s-t)} ds. \end{aligned} \quad (4.22)$$

Dropping the first two terms on the left-hand side of (4.22) and replacing ω by $\theta_{-t}\omega$, we obtain that, for all $t \geq T_1$,

$$\begin{aligned} & \int_{T_1}^t e^{\eta(s-t)} \|\nabla \tilde{u}(s, \theta_{-t}\omega, \tilde{u}_0(\theta_{-t}\omega))\|^2 ds + \alpha_1 \int_{T_1}^t e^{\eta(s-t)} \|u(s, \theta_{-t}\omega, u_0(\theta_{-t}\omega))\|_p^p ds \\ & \leq e^{\eta(T_1-t)} (\beta \|\tilde{u}(T_1, \theta_{-t}\omega, \tilde{u}_0(\theta_{-t}\omega))\|^2 + \alpha \|\tilde{v}(T_1, \theta_{-t}\omega, \tilde{v}_0(\theta_{-t}\omega))\|^2) \\ & + \int_{T_1}^t e^{\eta(s-t)} p_1(\theta_{s-t}\omega) ds + c \int_{T_1}^t e^{\eta(s-t)} ds. \end{aligned}$$

$$\begin{aligned} &\leq e^{\eta(T_1-t)} \left(\beta \|\tilde{u}(T_1, \theta_{-t}\omega, \tilde{u}_0(\theta_{-t}\omega))\|^2 + \alpha \|\tilde{v}(T_1, \theta_{-t}\omega, \tilde{v}_0(\theta_{-t}\omega))\|^2 \right) \\ &\quad + \int_{T_1-t}^0 e^{\eta\tau} p_1(\theta_\tau\omega) d\tau + \frac{c}{\eta} \end{aligned} \quad (4.23)$$

By (4.10), the second term on the right-hand side of (4.23) satisfies, for $t \geq T_1$,

$$\int_{T_1-t}^0 e^{\eta\tau} p_1(\theta_\tau\omega) d\tau \leq c_6 r(\omega) \int_{T_1-t}^0 e^{\frac{1}{2}\eta\tau} d\tau \leq \frac{2}{\eta} c_6 r(\omega). \quad (4.24)$$

Then, using (4.21) and (4.24), it follows from (4.23) that

$$\begin{aligned} &\int_{T_1}^t e^{\eta(s-t)} \|\nabla \tilde{u}(s, \theta_{-t}\omega, \tilde{u}_0(\theta_{-t}\omega))\|^2 ds + \alpha_1 \int_{T_1}^t e^{\eta(s-t)} \|u(s, \theta_{-t}\omega, u_0(\theta_{-t}\omega))\|_p^p ds \\ &\leq e^{-\eta t} \left(\beta \|\tilde{u}_0(\theta_{-t}\omega)\|^2 + \alpha \|\tilde{v}_0(\theta_{-t}\omega)\|^2 \right) + c(1 + r(\omega)). \end{aligned}$$

This completes the proof. $\square$

As a special case of Lemma 4.2, we have the following uniform estimates.

Lemma 4.3. *Assume that $g, h \in L^2(\mathbb{R}^n)$ and (3.4)-(3.7) hold. Let $B = \{B(\omega)\}_{\omega \in \Omega} \in \mathcal{D}$. Then for P -a.e. $\omega \in \Omega$, there exists $T_B(\omega) > 0$ such that for all $t \geq T_B(\omega)$,*

$$\begin{aligned} &\int_t^{t+1} \|u(s, \theta_{-t-1}\omega, u_0(\theta_{-t-1}\omega))\|_p^p ds \leq c(1 + r(\omega)), \\ &\int_t^{t+1} \|\nabla \tilde{u}(s, \theta_{-t-1}\omega, \tilde{u}_0(\theta_{-t-1}\omega))\|^2 ds \leq c(1 + r(\omega)), \end{aligned}$$

where c is a positive deterministic constant and $r(\omega)$ is the tempered function in (3.10).

Proof. First replacing t by $t+1$ and then replacing T_1 by t in (4.18), we find that

$$\begin{aligned} &\int_t^{t+1} e^{\eta(s-t-1)} \|\nabla \tilde{u}(s, \theta_{-t-1}\omega, \tilde{u}_0(\theta_{-t-1}\omega))\|^2 ds \\ &\leq ce^{-\eta(t+1)} \left(\|\tilde{u}_0(\theta_{-t-1}\omega)\|^2 + \|\tilde{v}_0(\theta_{-t-1}\omega)\|^2 \right) + c(1 + r(\omega)). \end{aligned} \quad (4.25)$$

Note that $e^{\eta(s-t-1)} \geq e^{-\eta}$ for $s \in [t, t+1]$. Hence from (4.25) we get that

$$\begin{aligned} &e^{-\eta} \int_t^{t+1} \|\nabla \tilde{u}(s, \theta_{-t-1}\omega, \tilde{u}_0(\theta_{-t-1}\omega))\|^2 ds \\ &\leq ce^{-\eta(t+1)} \left(\|\tilde{u}_0(\theta_{-t-1}\omega)\|^2 + \|\tilde{v}_0(\theta_{-t-1}\omega)\|^2 \right) + c(1 + r(\omega)). \end{aligned}$$

$$\begin{aligned}
&\leq ce^{-\eta(t+1)} (\|u_0(\theta_{-t-1}\omega)\|^2 + \|v_0(\theta_{-t-1}\omega)\|^2) \\
&+ ce^{-\eta(t+1)} (\|z_1(\theta_{-t-1}\omega)\|^2 + \|z_2(\theta_{-t-1}\omega)\|^2) + c(1+r(\omega)).
\end{aligned} \tag{4.26}$$

Since $\|u_0(\omega)\|^2$, $\|v_0(\omega)\|^2$, $\|z_1(\omega)\|^2$ and $\|z_2(\omega)\|^2$ are tempered, there is $T_B(\omega) > 0$ such that for all $t \geq T_B(\omega)$,

$$ce^{-\eta(t+1)} (\|u_0(\theta_{-t-1}\omega)\|^2 + \|v_0(\theta_{-t-1}\omega)\|^2 + \|z_1(\theta_{-t-1}\omega)\|^2 + \|z_2(\theta_{-t-1}\omega)\|^2) \leq c(1+r(\omega)),$$

which along with (4.26) shows that, for all $t \geq T_B(\omega)$,

$$\int_t^{t+1} \|\nabla \tilde{u}(s, \theta_{-t-1}\omega, \tilde{u}_0(\theta_{-t-1}\omega))\|^2 ds \leq 2e^\eta(1+r(\omega)). \tag{4.27}$$

Using (4.17) and repeating the above process, we can also find that, for $t \geq T_B(\omega)$,

$$\int_t^{t+1} \|u(s, \theta_{-t-1}\omega, u_0(\theta_{-t-1}\omega))\|_p^p ds \leq 2e^\eta(1+r(\omega)). \tag{4.28}$$

Then the lemma follows from (4.27)-(4.28). $\square$

Lemma 4.4. *Assume that $g, h \in L^2(\mathbb{R}^n)$ and (3.4)-(3.7) hold. Let $B = \{B(\omega)\}_{\omega \in \Omega} \in \mathcal{D}$. Then for P -a.e. $\omega \in \Omega$, there exists $T_B(\omega) > 0$ such that for all $t \geq T_B(\omega)$,*

$$\int_t^{t+1} \|\nabla u(s, \theta_{-t-1}\omega, u_0(\theta_{-t-1}\omega))\|^2 ds \leq c(1+r(\omega)),$$

where c is a positive deterministic constant and $r(\omega)$ is the tempered function in (3.10).

Proof. Let $T_B(\omega)$ be the positive constant in Lemma 4.3, take $t \geq T_B(\omega)$ and $s \in (t, t+1)$. By (3.15) we find that

$$\begin{aligned}
\|\nabla u(s, \theta_{-t-1}\omega, u_0(\theta_{-t-1}\omega))\|^2 &= \|\nabla \tilde{u}(s, \theta_{-t-1}\omega, \tilde{u}_0(\theta_{-t-1}\omega)) + \nabla z_1(\theta_{s-t-1}\omega)\|^2 \\
&\leq 2\|\nabla \tilde{u}(s, \theta_{-t-1}\omega, \tilde{u}_0(\theta_{-t-1}\omega))\|^2 + 2\|\nabla z_1(\theta_{s-t-1}\omega)\|^2.
\end{aligned} \tag{4.29}$$

By (3.12) we have

$$2\|\nabla z_1(\theta_{s-t-1}\omega)\|^2 = 2\|\nabla \phi_1\|^2 |y_1(\theta_{s-t-1}\omega_1)|^2 \leq ce^{\frac{\eta}{2}(t+1-s)} r(\omega) \leq ce^{\frac{\eta}{2}} r(\omega). \tag{4.30}$$

Now integrating (4.29) with respect to s over $(t, t+1)$, by Lemma 4.3 and inequality (4.30), we get that

$$\int_t^{t+1} \|\nabla u(s, \theta_{-t-1}\omega, u_0(\theta_{-t-1}\omega))\|^2 ds \leq c_1 + c_2 r(\omega). \tag{4.31}$$

Then the lemma follows from (4.31). $\square$

Next, we derive uniform estimates on u in $H^1(\mathbb{R}^n)$.

Lemma 4.5. *Assume that $g, h \in L^2(\mathbb{R}^n)$ and (3.4)-(3.7) hold. Let $B = \{B(\omega)\}_{\omega \in \Omega} \in \mathcal{D}$. Then for P -a.e. $\omega \in \Omega$, there exists $T_B(\omega) > 0$ such that for all $t \geq T_B(\omega)$,*

$$\|\nabla u(t, \theta_{-t}\omega, u_0(\theta_{-t}\omega))\|^2 \leq c(1 + r(\omega)),$$

where c is a positive deterministic constant and $r(\omega)$ is the tempered function in (3.10).

Proof. Taking the inner product of (3.13) with $\Delta \tilde{u}$ in $L^2(\mathbb{R}^n)$, we get that

$$\begin{aligned} & \frac{1}{2} \frac{d}{dt} \|\nabla \tilde{u}\|^2 + \lambda \|\nabla \tilde{u}\|^2 + \|\Delta \tilde{u}\|^2 \\ &= \alpha(\tilde{v}, \Delta \tilde{u}) - \int_{\mathbb{R}^n} f(x, u) \Delta \tilde{u} dx - (g + \Delta z_1(\theta_t \omega) - \alpha z_2(\theta_t \omega), \Delta \tilde{u}). \end{aligned} \quad (4.32)$$

Note that the first term on the right-hand side of (4.32) is bounded by

$$\alpha|(\tilde{v}, \Delta \tilde{u})| \leq \alpha \|\tilde{v}\| \|\Delta \tilde{u}\| \leq \frac{1}{4} \|\Delta \tilde{u}\|^2 + \alpha^2 \|\tilde{v}\|^2. \quad (4.33)$$

For the nonlinear term in (4.32), by (3.5)-(3.7), we have

$$\begin{aligned} & - \int_{\mathbb{R}^n} f(x, u) \Delta \tilde{u} dx = - \int_{\mathbb{R}^n} f(x, u) \Delta u dx + \int_{\mathbb{R}^n} f(x, u) \Delta z_1(\theta_t \omega) dx \\ &= \int_{\mathbb{R}^n} \frac{\partial f}{\partial x}(x, u) \nabla u dx + \int_{\mathbb{R}^n} \frac{\partial f}{\partial u}(x, u) |\nabla u|^2 dx + \int_{\mathbb{R}^n} f(x, u) \Delta z_1(\theta_t \omega) dx \\ &\leq \|\psi_3\| \|\nabla u\| + \beta \|\nabla u\|^2 + \int_{\mathbb{R}^n} |f(x, u)| |\Delta z_1(\theta_t \omega)| dx \\ &\leq \|\psi_3\| \|\nabla u\| + \beta \|\nabla u\|^2 + \alpha_2 \int_{\mathbb{R}^n} |u|^{p-1} |\Delta z_1(\theta_t \omega)| dx + \int_{\mathbb{R}^n} |\psi_2(x)| |\Delta z_1(\theta_t \omega)| dx \\ &\leq c \|\nabla u\|^2 + \frac{\alpha_2}{q} \int_{\mathbb{R}^n} |u|^p dx + \frac{\alpha_2}{p} \int_{\mathbb{R}^n} |\Delta z_1(\theta_t \omega)|^p dx + c(\|\psi_2\|^2 + \|\psi_3\|^2) + c \|\Delta z_1(\theta_t \omega)\|^2 \\ &\leq c(\|\nabla u\|^2 + \|u\|_p^p) + c(\|\Delta z_1(\theta_t \omega)\|^2 + \|\Delta z_1(\theta_t \omega)\|_p^p + 1). \end{aligned} \quad (4.34)$$

On the other hand, the last term on the right-hand side of (4.32) is bounded by

$$\begin{aligned} & |(g, \Delta \tilde{u})| + |(\Delta z_1(\theta_t \omega), \Delta \tilde{u})| + \alpha |(z_2(\theta_t \omega), \Delta \tilde{u})| \\ &\leq \frac{1}{4} \|\Delta \tilde{u}\|^2 + c(\|g\|^2 + \|\Delta z_1(\theta_t \omega)\|^2 + \|z_2(\theta_t \omega)\|^2). \end{aligned} \quad (4.35)$$

By (4.32)-(4.35) we get that

$$\begin{aligned} & \frac{d}{dt} \|\nabla \tilde{u}\|^2 + 2\lambda \|\nabla \tilde{u}\|^2 + \|\Delta \tilde{u}\|^2 \\ & \leq 2\alpha^2 \|\tilde{v}\|^2 + c (\|\nabla u\|^2 + \|u\|_p^p) + c (\|\Delta z_1(\theta_t \omega)\|^2 + \|\Delta z_1(\theta_t \omega)\|_p^p + \|z_2(\theta_t \omega)\|^2 + 1). \end{aligned} \quad (4.36)$$

Let

$$p_2(\theta_t \omega) = c (\|\Delta z_1(\theta_t \omega)\|^2 + \|\Delta z_1(\theta_t \omega)\|_p^p + \|z_2(\theta_t \omega)\|^2 + 1). \quad (4.37)$$

Since $z_1(\theta_t \omega) = \phi_1 y_1(\theta_t \omega_1)$ and $z_2(\theta_t \omega) = \phi_2 y_2(\theta_t \omega_2)$ with $\phi_1 \in H^2(\mathbb{R}^n) \cap W^{2,p}(\mathbb{R}^n)$ and $\phi_2 \in H^1(\mathbb{R}^n)$, we find that there are positive constants c_1 and c_2 such that

$$p_2(\theta_t \omega) \leq c_1 \sum_{j=1}^2 (|y_j(\theta_t \omega_j)|^2 + |y_j(\theta_t \omega_j)|_p^p) + c_2,$$

which along with (3.12) shows that

$$p_2(\theta_t \omega) \leq c_1 e^{\frac{\alpha}{2}|t|} r(\omega) + c_2, \quad \forall t \in \mathbb{R}. \quad (4.38)$$

By (4.36)-(4.37), we find that

$$\frac{d}{dt} \|\nabla \tilde{u}\|^2 \leq c (\|\nabla u\|^2 + \|u\|_p^p + \|\tilde{v}\|^2) + p_2(\theta_t \omega). \quad (4.39)$$

Let $T_B(\omega)$ be the positive constant in Lemma 4.3, take $t \geq T_B(\omega)$ and $s \in (t, t+1)$. Then integrate (4.39) over $(s, t+1)$ to get

$$\begin{aligned} & \|\nabla \tilde{u}(t+1, \omega, \tilde{u}_0(\omega))\|^2 \leq \|\nabla \tilde{u}(s, \omega, \tilde{u}_0(\omega))\|^2 + \int_s^{t+1} p_2(\theta_\tau \omega) d\tau \\ & + c \int_s^{t+1} (\|\nabla u(\tau, \omega, u_0(\omega))\|^2 + \|u(\tau, \omega, u_0(\omega))\|_p^p + \|\tilde{v}(\tau, \omega, \tilde{v}_0(\omega))\|^2) d\tau \\ & \leq \|\nabla \tilde{u}(s, \omega, \tilde{u}_0(\omega))\|^2 + \int_t^{t+1} p_2(\theta_\tau \omega) d\tau \\ & + c \int_t^{t+1} (\|\nabla u(\tau, \omega, u_0(\omega))\|^2 + \|u(\tau, \omega, u_0(\omega))\|_p^p + \|\tilde{v}(\tau, \omega, \tilde{v}_0(\omega))\|^2) d\tau. \end{aligned}$$

Now integrating the above with respect to s over $(t, t+1)$, we find that

$$\begin{aligned} & \|\nabla \tilde{u}(t+1, \omega, \tilde{u}_0(\omega))\|^2 \leq \int_t^{t+1} \|\nabla \tilde{u}(s, \omega, \tilde{u}_0(\omega))\|^2 ds + \int_t^{t+1} p_2(\theta_\tau \omega) d\tau \\ & + c \int_t^{t+1} (\|\nabla u(\tau, \omega, u_0(\omega))\|^2 + \|u(\tau, \omega, u_0(\omega))\|_p^p + \|\tilde{v}(\tau, \omega, \tilde{v}_0(\omega))\|^2) d\tau. \end{aligned}$$

Replacing ω by $\theta_{-t-1}\omega$, we obtain that

$$\begin{aligned}
& \|\nabla \tilde{u}(t+1, \theta_{-t-1}\omega, \tilde{u}_0(\theta_{-t-1}\omega))\|^2 \\
& \leq \int_t^{t+1} \|\nabla \tilde{u}(s, \theta_{-t-1}\omega, \tilde{u}_0(\theta_{-t-1}\omega))\|^2 ds + \int_t^{t+1} p_2(\theta_{\tau-t-1}\omega) d\tau \\
& + c \int_t^{t+1} (\|\nabla u(\tau, \theta_{-t-1}\omega, u_0(\theta_{-t-1}\omega))\|^2 + \|u(\tau, \theta_{-t-1}\omega, u_0(\theta_{-t-1}\omega))\|_p^p) d\tau \\
& + c \int_t^{t+1} \|\tilde{v}(\tau, \theta_{-t-1}\omega, \tilde{v}_0(\theta_{-t-1}\omega))\|^2 d\tau.
\end{aligned} \tag{4.40}$$

By Lemmas 4.3 and 4.4, it follows from (4.40) and (4.38) that, for all $t \geq T_B(\omega)$,

$$\begin{aligned}
\|\nabla \tilde{u}(t+1, \theta_{-t-1}\omega, \tilde{u}_0(\theta_{-t-1}\omega))\|^2 & \leq c_3 + c_4 r(\omega) + \int_{-1}^0 p_2(\theta_s \omega) ds \\
& \leq c_3 + c_4 r(\omega) + \int_{-1}^0 \left(c_1 e^{-\frac{\eta}{2}s} r(\omega) + c_2 \right) ds \leq c_5 + c_6 r(\omega).
\end{aligned} \tag{4.41}$$

Then by (4.41) and (3.15), we have, for all $t \geq T_B(\omega)$,

$$\begin{aligned}
\|\nabla u(t+1, \theta_{-t-1}\omega, u_0(\theta_{-t-1}\omega))\|^2 & = \|\nabla \tilde{u}(t+1, \theta_{-t-1}\omega, \tilde{u}_0(\theta_{-t-1}\omega)) + \nabla z_1(\omega)\|^2 \\
& \leq 2\|\nabla \tilde{u}(t+1, \theta_{-t-1}\omega, \tilde{u}_0(\theta_{-t-1}\omega))\|^2 + 2\|\nabla z_1(\omega)\|^2 \leq c_7 + c_8 r(\omega),
\end{aligned}$$

which completes the proof. $\square$

Lemma 4.6. *Assume that $g, h \in L^2(\mathbb{R}^n)$ and (3.4)-(3.7) hold. Let $B = \{B(\omega)\}_{\omega \in \Omega} \in \mathcal{D}$. Then for every $\epsilon > 0$ and P -a.e. $\omega \in \Omega$, there exist $T = T_B(\omega, \epsilon) > 0$ and $R = R(\omega, \epsilon) > 0$ such that for all $t \geq T$,*

$$\int_{|x| \geq R} (|\tilde{u}(t, \theta_{-t}\omega, \tilde{u}_0(\theta_{-t}\omega))|^2 + |\tilde{v}(t, \theta_{-t}\omega, \tilde{v}_0(\theta_{-t}\omega))|^2) dx \leq \epsilon.$$

Proof. Let ρ be a smooth function defined on $\mathbb{R}^+$ such that $0 \leq \rho(s) \leq 1$ for all $s \in \mathbb{R}^+$, and

$$\rho(s) = \begin{cases} 0 & \text{for } 0 \leq s \leq 1; \\ 1 & \text{for } s \geq 2. \end{cases}$$

Then there exists a positive constant c such that $|\rho'(s)| \leq c$ for all $s \in \mathbb{R}^+$.

Taking the inner product of (3.13) with $\beta \rho(\frac{|x|^2}{k^2}) \tilde{u}$ in $L^2(\mathbb{R}^n)$ we find that

$$\frac{1}{2} \beta \frac{d}{dt} \int_{\mathbb{R}^n} \rho\left(\frac{|x|^2}{k^2}\right) |\tilde{u}|^2 dx + \lambda \beta \int_{\mathbb{R}^n} \rho\left(\frac{|x|^2}{k^2}\right) |\tilde{u}|^2 dx - \beta \int_{\mathbb{R}^n} \rho\left(\frac{|x|^2}{k^2}\right) \tilde{u} \Delta \tilde{u} dx + \alpha \beta \int_{\mathbb{R}^n} \rho\left(\frac{|x|^2}{k^2}\right) \tilde{u} \tilde{v} dx$$

$$= \beta \int_{\mathbb{R}^n} f(x, u) \rho\left(\frac{|x|^2}{k^2}\right) \tilde{u} dx + \beta \int_{\mathbb{R}^n} (g + \Delta z_1(\theta_t \omega) - \alpha z_2(\theta_t \omega)) \rho\left(\frac{|x|^2}{k^2}\right) \tilde{u} dx. \quad (4.42)$$

Taking the inner product of (3.14) with $\alpha \rho\left(\frac{|x|^2}{k^2}\right) \tilde{v}$ in $L^2(\mathbb{R}^n)$ we find that

$$\begin{aligned} & \frac{1}{2} \alpha \frac{d}{dt} \int_{\mathbb{R}^n} \rho\left(\frac{|x|^2}{k^2}\right) |\tilde{v}|^2 dx + \alpha \delta \int_{\mathbb{R}^n} \rho\left(\frac{|x|^2}{k^2}\right) |\tilde{v}|^2 dx - \alpha \beta \int_{\mathbb{R}^n} \rho\left(\frac{|x|^2}{k^2}\right) \tilde{u} \tilde{v} dx \\ &= \alpha \int_{\mathbb{R}^n} \rho\left(\frac{|x|^2}{k^2}\right) h \tilde{v} dx + \alpha \beta \int_{\mathbb{R}^n} \rho\left(\frac{|x|^2}{k^2}\right) z_1(\theta_t \omega) \tilde{v} dx. \end{aligned} \quad (4.43)$$

Adding (4.2) and (4.3), we obtain that

$$\begin{aligned} & \frac{1}{2} \frac{d}{dt} \left(\beta \int_{\mathbb{R}^n} \rho\left(\frac{|x|^2}{k^2}\right) |\tilde{u}|^2 dx + \alpha \int_{\mathbb{R}^n} \rho\left(\frac{|x|^2}{k^2}\right) |\tilde{v}|^2 dx \right) \\ &+ \lambda \beta \int_{\mathbb{R}^n} \rho\left(\frac{|x|^2}{k^2}\right) |\tilde{u}|^2 dx + \alpha \delta \int_{\mathbb{R}^n} \rho\left(\frac{|x|^2}{k^2}\right) |\tilde{v}|^2 dx - \beta \int_{\mathbb{R}^n} \rho\left(\frac{|x|^2}{k^2}\right) \tilde{u} \Delta \tilde{u} dx \\ &= \beta \int_{\mathbb{R}^n} f(x, u) \rho\left(\frac{|x|^2}{k^2}\right) \tilde{u} dx + \beta \int_{\mathbb{R}^n} (g + \Delta z_1(\theta_t \omega) - \alpha z_2(\theta_t \omega)) \rho\left(\frac{|x|^2}{k^2}\right) \tilde{u} dx \\ &+ \alpha \int_{\mathbb{R}^n} \rho\left(\frac{|x|^2}{k^2}\right) h \tilde{v} dx + \alpha \beta \int_{\mathbb{R}^n} \rho\left(\frac{|x|^2}{k^2}\right) z_1(\theta_t \omega) \tilde{v} dx. \end{aligned} \quad (4.44)$$

We now estimate the terms in (4.44) as follows. First we have

$$\begin{aligned} -\beta \int_{\mathbb{R}^n} \rho\left(\frac{|x|^2}{k^2}\right) \tilde{u} \Delta \tilde{u} dx &= \beta \int_{\mathbb{R}^n} |\nabla \tilde{u}|^2 \rho\left(\frac{|x|^2}{k^2}\right) dx + \beta \int_{\mathbb{R}^n} \tilde{u} \rho'\left(\frac{|x|^2}{k^2}\right) \frac{2x}{k^2} \cdot \nabla \tilde{u} dx \\ &= \beta \int_{\mathbb{R}^n} |\nabla \tilde{u}|^2 \rho\left(\frac{|x|^2}{k^2}\right) dx + \beta \int_{k \leq |x| \leq \sqrt{2}k} \tilde{u} \rho'\left(\frac{|x|^2}{k^2}\right) \frac{2x}{k^2} \cdot \nabla \tilde{u} dx. \end{aligned} \quad (4.45)$$

Note that the second term on the right-hand side of (4.45) is bounded by

$$\begin{aligned} \beta \left| \int_{k \leq |x| \leq \sqrt{2}k} \tilde{u} \rho'\left(\frac{|x|^2}{k^2}\right) \frac{2x}{k^2} \cdot \nabla \tilde{u} dx \right| &\leq \frac{2\sqrt{2}}{k} \beta \int_{k \leq |x| \leq \sqrt{2}k} |\tilde{u}| \left| \rho'\left(\frac{|x|^2}{k^2}\right) \right| |\nabla \tilde{u}| dx \\ &\leq \frac{c}{k} \int_{\mathbb{R}^n} |\tilde{u}| |\nabla \tilde{u}| dx \leq \frac{c}{k} (\|\tilde{u}\|^2 + \|\nabla \tilde{u}\|^2). \end{aligned} \quad (4.46)$$

By (4.45)-(4.46), we find that

$$-\beta \int_{\mathbb{R}^n} \rho\left(\frac{|x|^2}{k^2}\right) \tilde{u} \Delta \tilde{u} dx \geq \beta \int_{\mathbb{R}^n} |\nabla \tilde{u}|^2 \rho\left(\frac{|x|^2}{k^2}\right) dx - \frac{c}{k} (\|\tilde{u}\|^2 + \|\nabla \tilde{u}\|^2). \quad (4.47)$$

For the nonlinear term in (4.44), we have

$$\beta \int_{\mathbb{R}^n} f(x, u) \rho\left(\frac{|x|^2}{k^2}\right) \tilde{u} dx = \beta \int_{\mathbb{R}^n} f(x, u) \rho\left(\frac{|x|^2}{k^2}\right) u dx - \beta \int_{\mathbb{R}^n} f(x, u) \rho\left(\frac{|x|^2}{k^2}\right) z_1(\theta_t \omega) dx. \quad (4.48)$$

By (3.4), the first term on the right-hand side of (4.48) is bounded by

$$\beta \int_{\mathbb{R}^n} f(x, u) \rho\left(\frac{|x|^2}{k^2}\right) u dx \leq -\alpha_1 \beta \int_{\mathbb{R}^n} |u|^p \rho\left(\frac{|x|^2}{k^2}\right) dx + \beta \int_{\mathbb{R}^n} \psi_1 \rho\left(\frac{|x|^2}{k^2}\right) dx. \quad (4.49)$$

By (3.5), the second term on the right-hand side of (4.48) is bounded by

$$\begin{aligned} & \beta \left| \int_{\mathbb{R}^n} f(x, u) \rho\left(\frac{|x|^2}{k^2}\right) z_1(\theta_t \omega) dx \right| \\ & \leq \alpha_2 \beta \int_{\mathbb{R}^n} |u|^{p-1} \rho\left(\frac{|x|^2}{k^2}\right) |z_1(\theta_t \omega)| dx + \beta \int_{\mathbb{R}^n} |\psi_2| \rho\left(\frac{|x|^2}{k^2}\right) |z_1(\theta_t \omega)| dx \\ & \leq \frac{1}{2} \alpha_1 \beta \int_{\mathbb{R}^n} |u|^p \rho\left(\frac{|x|^2}{k^2}\right) dx + c \int_{\mathbb{R}^n} |z_1(\theta_t \omega)|^p \rho\left(\frac{|x|^2}{k^2}\right) dx \\ & \quad + \frac{1}{2} \beta \int_{\mathbb{R}^n} |z_1(\theta_t \omega)|^2 \rho\left(\frac{|x|^2}{k^2}\right) dx + \frac{1}{2} \beta \int_{\mathbb{R}^n} \psi_2^2 \rho\left(\frac{|x|^2}{k^2}\right) dx. \end{aligned} \quad (4.50)$$

Then it follows from (4.48)-(4.50) that

$$\begin{aligned} \beta \int_{\mathbb{R}^n} f(x, u) \rho\left(\frac{|x|^2}{k^2}\right) \tilde{u} dx & \leq -\frac{1}{2} \alpha_1 \beta \int_{\mathbb{R}^n} |u|^p \rho\left(\frac{|x|^2}{k^2}\right) dx + \beta \int_{\mathbb{R}^n} \psi_1 \rho\left(\frac{|x|^2}{k^2}\right) dx \\ & \quad + \frac{1}{2} \beta \int_{\mathbb{R}^n} \psi_2^2 \rho\left(\frac{|x|^2}{k^2}\right) dx + c \int_{\mathbb{R}^n} (|z_1(\theta_t \omega)|^p + |z_1(\theta_t \omega)|^2) \rho\left(\frac{|x|^2}{k^2}\right) dx. \end{aligned} \quad (4.51)$$

For the second term on the right-hand side of (4.44), we have that

$$\begin{aligned} & \beta \left| \int_{\mathbb{R}^n} (g + \Delta z_1(\theta_t \omega) - \alpha z_2(\theta_t \omega)) \rho\left(\frac{|x|^2}{k^2}\right) \tilde{u} dx \right| \\ & \leq \frac{1}{2} \lambda \beta \int_{\mathbb{R}^n} \rho\left(\frac{|x|^2}{k^2}\right) |\tilde{u}|^2 dx + c \int_{\mathbb{R}^n} (g^2 + |\Delta z_1(\theta_t \omega)|^2 + |z_2(\theta_t \omega)|^2) \rho\left(\frac{|x|^2}{k^2}\right) dx. \end{aligned} \quad (4.52)$$

For the last two terms on the right-hand side of (4.44), we find that

$$\begin{aligned} & \alpha \int_{\mathbb{R}^n} \rho\left(\frac{|x|^2}{k^2}\right) h \tilde{v} dx + \alpha \beta \int_{\mathbb{R}^n} \rho\left(\frac{|x|^2}{k^2}\right) z_1(\theta_t \omega) \tilde{v} dx \\ & \leq \frac{1}{2} \alpha \delta \int_{\mathbb{R}^n} \rho\left(\frac{|x|^2}{k^2}\right) |\tilde{v}|^2 dx + c \int_{\mathbb{R}^n} \rho\left(\frac{|x|^2}{k^2}\right) (|z_1(\theta_t \omega)|^2 + |h|^2) dx. \end{aligned} \quad (4.53)$$

Finally, by (4.44), (4.47) and (4.51)-(4.53), we obtain that

$$\begin{aligned} & \frac{1}{2} \frac{d}{dt} \left(\beta \int_{\mathbb{R}^n} \rho\left(\frac{|x|^2}{k^2}\right) |\tilde{u}|^2 dx + \alpha \int_{\mathbb{R}^n} \rho\left(\frac{|x|^2}{k^2}\right) |\tilde{v}|^2 dx \right) \\ & \quad + \frac{1}{2} \lambda \beta \int_{\mathbb{R}^n} \rho\left(\frac{|x|^2}{k^2}\right) |\tilde{u}|^2 dx + \frac{1}{2} \alpha \delta \int_{\mathbb{R}^n} \rho\left(\frac{|x|^2}{k^2}\right) |\tilde{v}|^2 dx \end{aligned}$$

$$\begin{aligned}
& + \frac{1}{2} \alpha_1 \beta \int_{\mathbb{R}^n} \rho\left(\frac{|x|^2}{k^2}\right) |u|^p dx + \beta \int_{\mathbb{R}^n} \rho\left(\frac{|x|^2}{k^2}\right) |\nabla \tilde{u}|^2 dx \\
& \leq \frac{c}{k} (\|\nabla \tilde{u}\|^2 + \|\tilde{u}\|^2) + c \int_{\mathbb{R}^n} (|\psi_1| + |\psi_2|^2 + g^2 + h^2) \rho\left(\frac{|x|^2}{k^2}\right) dx \\
& + c \int_{\mathbb{R}^n} (|\Delta z_1(\theta_t \omega)|^2 + |z_1(\theta_t \omega)|^2 + |z_1(\theta_t \omega)|^p + |z_2(\theta_t \omega)|^2) \rho\left(\frac{|x|^2}{k^2}\right) dx.
\end{aligned} \tag{4.54}$$

Note that (4.54) implies that

$$\begin{aligned}
& \frac{d}{dt} \left(\beta \int_{\mathbb{R}^n} \rho\left(\frac{|x|^2}{k^2}\right) |\tilde{u}|^2 dx + \alpha \int_{\mathbb{R}^n} \rho\left(\frac{|x|^2}{k^2}\right) |\tilde{v}|^2 dx \right) \\
& + \eta \left(\beta \int_{\mathbb{R}^n} \rho\left(\frac{|x|^2}{k^2}\right) |\tilde{u}|^2 dx + \alpha \int_{\mathbb{R}^n} \rho\left(\frac{|x|^2}{k^2}\right) |\tilde{v}|^2 dx \right) \\
& \leq \frac{c}{k} (\|\nabla \tilde{u}\|^2 + \|\tilde{u}\|^2) + c \int_{\mathbb{R}^n} (|\psi_1| + |\psi_2|^2 + g^2 + h^2) \rho\left(\frac{|x|^2}{k^2}\right) dx \\
& + c \int_{\mathbb{R}^n} (|\Delta z_1(\theta_t \omega)|^2 + |z_1(\theta_t \omega)|^2 + |z_1(\theta_t \omega)|^p + |z_2(\theta_t \omega)|^2) \rho\left(\frac{|x|^2}{k^2}\right) dx.
\end{aligned} \tag{4.55}$$

By Lemmas 4.1 and 4.5, there is $T_1 = T_1(B, \omega) > 0$ such that for all $t \geq T_1$,

$$\|\tilde{u}(t, \theta_{-t} \omega, \tilde{u}_0(\theta_{-t} \omega))\|_{H^1(\mathbb{R}^n)}^2 \leq c(1 + r(\omega)). \tag{4.56}$$

Now integrating (4.55) over (T_1, t) , we get that, for all $t \geq T_1$,

$$\begin{aligned}
& \beta \int_{\mathbb{R}^n} \rho\left(\frac{|x|^2}{k^2}\right) |\tilde{u}(t, \omega, \tilde{u}_0(\omega))|^2 dx + \alpha \int_{\mathbb{R}^n} \rho\left(\frac{|x|^2}{k^2}\right) |\tilde{v}(t, \omega, \tilde{v}_0(\omega))|^2 dx \\
& \leq e^{\eta(T_1-t)} \left(\beta \int_{\mathbb{R}^n} \rho\left(\frac{|x|^2}{k^2}\right) |\tilde{u}(T_1, \omega, \tilde{u}_0(\omega))|^2 dx + \alpha \int_{\mathbb{R}^n} \rho\left(\frac{|x|^2}{k^2}\right) |\tilde{v}(T_1, \omega, \tilde{v}_0(\omega))|^2 dx \right) \\
& + \frac{c}{k} \int_{T_1}^t e^{\eta(s-t)} (\|\nabla \tilde{u}(s, \omega, \tilde{u}_0(\omega))\|^2 + \|\tilde{u}(s, \omega, \tilde{u}_0(\omega))\|^2) ds \\
& + c \int_{T_1}^t e^{\eta(s-t)} \int_{\mathbb{R}^n} (|\psi_1| + |\psi_2|^2 + g^2 + h^2) \rho\left(\frac{|x|^2}{k^2}\right) dx ds \\
& + c \int_{T_1}^t e^{\eta(s-t)} \int_{\mathbb{R}^n} (|\Delta z_1(\theta_s \omega)|^2 + |z_1(\theta_s \omega)|^2 + |z_1(\theta_s \omega)|^p + |z_2(\theta_s \omega)|^2) \rho\left(\frac{|x|^2}{k^2}\right) dx ds.
\end{aligned} \tag{4.57}$$

Replacing ω by $\theta_{-t} \omega$, we obtain from (4.57) that, for all $t \geq T_1$,

$$\begin{aligned}
& \beta \int_{\mathbb{R}^n} \rho\left(\frac{|x|^2}{k^2}\right) |\tilde{u}(t, \theta_{-t} \omega, \tilde{u}_0(\theta_{-t} \omega))|^2 dx + \alpha \int_{\mathbb{R}^n} \rho\left(\frac{|x|^2}{k^2}\right) |\tilde{v}(t, \theta_{-t} \omega, \tilde{v}_0(\theta_{-t} \omega))|^2 dx \\
& \leq \beta e^{\eta(T_1-t)} \int_{\mathbb{R}^n} \rho\left(\frac{|x|^2}{k^2}\right) |\tilde{u}(T_1, \theta_{-t} \omega, \tilde{u}_0(\theta_{-t} \omega))|^2 dx
\end{aligned}$$

$$\begin{aligned}
& + \alpha e^{\eta(T_1-t)} \int_{\mathbb{R}^n} \rho\left(\frac{|x|^2}{k^2}\right) |\tilde{v}(T_1, \theta_{-t}\omega, \tilde{v}_0(\theta_{-t}\omega))|^2 dx \\
& + \frac{c}{k} \int_{T_1}^t e^{\eta(s-t)} \|\nabla \tilde{u}(s, \theta_{-t}\omega, \tilde{u}_0(\theta_{-t}\omega))\|^2 ds \\
& + \frac{c}{k} \int_{T_1}^t e^{\eta(s-t)} \|\tilde{u}(s, \theta_{-t}\omega, \tilde{u}_0(\theta_{-t}\omega))\|^2 ds \\
& + c \int_{T_1}^t e^{\eta(s-t)} \int_{\mathbb{R}^n} (|\psi_1| + |\psi_2|^2 + g^2 + h^2) \rho\left(\frac{|x|^2}{k^2}\right) dx ds \\
& + c \int_{T_1}^t e^{\eta(s-t)} \int_{\mathbb{R}^n} (|\Delta z_1(\theta_{s-t}\omega)|^2 + |z_1(\theta_{s-t}\omega)|^2 + |z_1(\theta_{s-t}\omega)|^p + |z_2(\theta_{s-t}\omega)|^2) \rho\left(\frac{|x|^2}{k^2}\right) dx ds. \quad (4.58)
\end{aligned}$$

In what follows, we estimate the terms in (4.58). First replacing t by T_1 and then replacing ω by $\theta_{-t}\omega$ in (4.13), we have the following bounds for the first two terms on the right-hand side of (4.58).

$$\begin{aligned}
& \beta e^{\eta(T_1-t)} \int_{\mathbb{R}^n} \rho\left(\frac{|x|^2}{k^2}\right) |\tilde{u}(T_1, \theta_{-t}\omega, \tilde{u}_0(\theta_{-t}\omega))|^2 dx \\
& + \alpha e^{\eta(T_1-t)} \int_{\mathbb{R}^n} \rho\left(\frac{|x|^2}{k^2}\right) |\tilde{v}(T_1, \theta_{-t}\omega, \tilde{v}_0(\theta_{-t}\omega))|^2 dx \\
& \leq e^{\eta(T_1-t)} \left(e^{-\eta T_1} (\beta \|\tilde{u}_0(\theta_{-t}\omega)\|^2 + \alpha \|\tilde{v}_0(\theta_{-t}\omega)\|^2) + \int_0^{T_1} e^{\eta(\tau-T_1)} p_1(\theta_{\tau-t}\omega) d\tau + c \right) \\
& \leq e^{-\eta t} (\beta \|\tilde{u}_0(\theta_{-t}\omega)\|^2 + \alpha \|\tilde{v}_0(\theta_{-t}\omega)\|^2) + \int_0^{T_1} e^{\eta(\tau-t)} p_1(\theta_{\tau-t}\omega) d\tau + c e^{\eta(T_1-t)} \\
& \leq e^{-\eta t} (\beta \|\tilde{u}_0(\theta_{-t}\omega)\|^2 + \alpha \|\tilde{v}_0(\theta_{-t}\omega)\|^2) + \int_{-t}^{T_1-t} e^{\eta\tau} p_1(\theta_\tau\omega) d\tau + c e^{\eta(T_1-t)} \\
& \leq e^{-\eta t} (\beta \|\tilde{u}_0(\theta_{-t}\omega)\|^2 + \alpha \|\tilde{v}_0(\theta_{-t}\omega)\|^2) + \int_{-t}^{T_1-t} e^{\frac{1}{2}\eta\tau} c_6 r(\omega) d\tau + c e^{\eta(T_1-t)} \\
& \leq e^{-\eta t} (\beta \|\tilde{u}_0(\theta_{-t}\omega)\|^2 + \alpha \|\tilde{v}_0(\theta_{-t}\omega)\|^2) + \frac{2}{\eta} c_6 r(\omega) e^{\frac{1}{2}\eta(T_1-t)} + c e^{\eta(T_1-t)} \quad (4.59)
\end{aligned}$$

where we have used (4.10). By (4.59), we find that, given $\epsilon > 0$, there is $T_2 = (B, \omega, \epsilon) > T_1$ such that for all $t \geq T_2$,

$$e^{\eta(T_1-t)} \left(\int_{\mathbb{R}^n} \rho\left(\frac{|x|^2}{k^2}\right) (\beta |\tilde{u}(T_1, \theta_{-t}\omega, \tilde{u}_0(\theta_{-t}\omega))|^2 + \alpha |\tilde{v}(T_1, \theta_{-t}\omega, \tilde{v}_0(\theta_{-t}\omega))|^2) dx \right) \leq \epsilon. \quad (4.60)$$

By Lemma 4.2, there is $T_3 = T_3(B, \omega) > T_1$ such that the third term on the right-hand side of (4.58) satisfies, for all $t \geq T_3$,

$$\frac{c}{k} \int_{T_1}^t e^{\eta(s-t)} \|\nabla \tilde{u}(s, \theta_{-t}\omega, \tilde{u}_0(\theta_{-t}\omega))\|^2 ds \leq \frac{c}{k} (1 + r(\omega)).$$

And hence, there is $R_1 = R_1(\omega, \epsilon) > 0$ such that for all $t \geq T_3$ and $k \geq R_1$,

$$\frac{c}{k} \int_{T_1}^t e^{\lambda(s-t)} \|\nabla v(s, \theta_{-t}\omega, v_0(\theta_{-t}\omega))\|^2 ds \leq \epsilon. \quad (4.61)$$

First replacing t by s and then replacing ω by $\theta_{-t}\omega$ in (4.13), we find that the fourth term on the right-hand side of (4.58) satisfies

$$\begin{aligned} & \frac{c}{k} \int_{T_1}^t e^{\eta(s-t)} \|\tilde{u}(s, \theta_{-t}\omega, \tilde{u}_0(\theta_{-t}\omega))\|^2 ds \\ & \leq \frac{c}{k} \int_{T_1}^t e^{-\eta t} (\beta \|\tilde{u}_0(\theta_{-t}\omega)\|^2 + \alpha \|\tilde{v}_0(\theta_{-t}\omega)\|^2) ds \\ & + \frac{c}{k} \int_{T_1}^t e^{\eta(s-t)} \int_0^s e^{\eta(\tau-s)} p_1(\theta_{\tau-t}\omega) d\tau ds + \frac{c}{k} \int_{T_1}^t e^{\lambda(s-t)} ds \\ & \leq \frac{c}{k} e^{-\eta t} (t - T_1) (\beta \|\tilde{u}_0(\theta_{-t}\omega)\|^2 + \alpha \|\tilde{v}_0(\theta_{-t}\omega)\|^2) \\ & \quad + \frac{c}{k} + \frac{c}{k} \int_{T_1}^t \int_0^s e^{\eta(\tau-t)} p_1(\theta_{\tau-t}\omega) d\tau ds \\ & \leq \frac{c}{k} e^{-\eta t} (t - T_1) (\beta \|\tilde{u}_0(\theta_{-t}\omega)\|^2 + \alpha \|\tilde{v}_0(\theta_{-t}\omega)\|^2) \\ & \quad + \frac{c}{k} + \frac{c}{k} \int_{T_1}^t \int_{-t}^{s-t} e^{\eta\tau} p_1(\theta_{\tau}\omega) d\tau ds \\ & \leq \frac{c}{k} e^{-\eta t} (t - T_1) (\beta \|\tilde{u}_0(\theta_{-t}\omega)\|^2 + \alpha \|\tilde{v}_0(\theta_{-t}\omega)\|^2) \\ & \quad + \frac{c}{k} + \frac{c}{k} r(\omega) \int_{T_1}^t \int_{-t}^{s-t} e^{\frac{1}{2}\eta\tau} d\tau ds \\ & \leq \frac{c}{k} e^{-\eta t} (t - T_1) (\beta \|\tilde{u}_0(\theta_{-t}\omega)\|^2 + \alpha \|\tilde{v}_0(\theta_{-t}\omega)\|^2) + \frac{c}{k} + \frac{4c}{\eta^2 k} r(\omega), \end{aligned}$$

which shows that, there is $T_4 = T_4(B, \omega, \epsilon) > T_1$ and $R_2 = R_2(\omega, \epsilon)$ such that for all $t \geq T_4$ and $k \geq R_2$,

$$\frac{c}{k} \int_{T_1}^t e^{\eta(s-t)} \|\tilde{u}(s, \theta_{-t}\omega, \tilde{u}_0(\theta_{-t}\omega))\|^2 ds \leq \epsilon. \quad (4.62)$$

Note that $\psi_1 \in L^1(\mathbb{R}^n)$ and $\psi_2, g, h \in L^2(\mathbb{R}^n)$. Therefore, there is $R_3 = R_3(\epsilon)$ such that for all $k \geq R_3$,

$$\int_{|x| \geq k} (|\psi_1| + |\psi_2|^2 + |g|^2 + |h|^2) dx \leq \epsilon.$$

Then for the fifth term on the right-hand side of (4.58), we have

$$c \int_{T_1}^t e^{\eta(s-t)} \int_{\mathbb{R}^n} (|\psi_1| + |\psi_2|^2 + |g|^2 + |h|^2) \rho\left(\frac{|x|^2}{k^2}\right) dx ds$$

$$\leq c \int_{T_1}^t e^{\eta(s-t)} \int_{|x| \geq k} (|\psi_1| + |\psi_2|^2 + |g|^2 + |h|^2) dx ds \leq c\epsilon \int_{T_1}^t e^{\eta(s-t)} ds \leq c\epsilon, \quad (4.63)$$

where c is independent of ϵ . Note that $\phi_1 \in H^2(\mathbb{R}^n) \cap W^{2,p}(\mathbb{R}^n)$ and $\phi_2 \in H^1(\mathbb{R}^n)$. Hence there is $R_4 = R_4(\omega, \epsilon)$ such that for all $k \geq R_4$,

$$\int_{|x| \geq k} (|\phi_1(x)|^2 + |\phi_1(x)|^p + |\Delta \phi_1(x)|^2 + |\phi_2(x)|^2) dx \leq \frac{\epsilon}{r(\omega)}, \quad (4.64)$$

where $r(\omega)$ is the tempered function in (3.10). By (4.64) and (3.10)-(3.11), we have the following bounds for the last term on the right-hand side of (4.58):

$$\begin{aligned} & c \int_{T_1}^t e^{\eta(s-t)} \int_{\mathbb{R}^n} (|\Delta z_1(\theta_{s-t}\omega)|^2 + |z_1(\theta_{s-t}\omega)|^2 + |z_1(\theta_{s-t}\omega)|^p + |z_2(\theta_{s-t}\omega)|^2) \rho\left(\frac{|x|^2}{k^2}\right) dx ds \\ & \leq c \int_{T_1}^t e^{\eta(s-t)} \int_{|x| \geq k} (|\Delta z_1(\theta_{s-t}\omega)|^2 + |z_1(\theta_{s-t}\omega)|^2 + |z_1(\theta_{s-t}\omega)|^p + |z_2(\theta_{s-t}\omega)|^2) dx ds \\ & \leq c \int_{T_1}^t e^{\eta(s-t)} \int_{|x| \geq k} (|\Delta \phi_1|^2 |y_1(\theta_{s-t}\omega_1)|^2 + |\phi_1|^2 |y_1(\theta_{s-t}\omega_1)|^2 + |\phi_1|^p |y_1(\theta_{s-t}\omega_1)|^p + |\phi_2|^2 |y_2(\theta_{s-t}\omega_2)|^2) dx ds \\ & \leq \frac{c\epsilon}{r(\omega)} \int_{T_1}^t e^{\eta(s-t)} \sum_{j=1}^2 (|y_j(\theta_{s-t}\omega_j)|^2 + |y_j(\theta_{s-t}\omega_j)|^p) ds \leq \frac{c\epsilon}{r(\omega)} \int_{T_1}^t e^{\eta(s-t)} r(\theta_{s-t}\omega) ds \\ & \leq \frac{c\epsilon}{r(\omega)} \int_{T_1-t}^0 e^{\eta\tau} r(\theta_\tau\omega) d\tau \leq \frac{c\epsilon}{r(\omega)} \int_{T_1-t}^0 e^{\frac{1}{2}\eta\tau} r(\omega) d\tau \leq c\epsilon. \end{aligned} \quad (4.65)$$

Let $T_5 = T_5(B, \omega, \epsilon) = \max\{T_1, T_2, T_3, T_4\}$ and $R_5 = R_5(\omega, \epsilon) = \max\{R_1, R_2, R_3, R_4\}$. Then it follows from (4.58), (4.60)-(4.65) that, for all $t \geq T_5$ and $k \geq R_5$,

$$\beta \int_{\mathbb{R}^n} \rho\left(\frac{|x|^2}{k^2}\right) |\tilde{u}(t, \theta_{-t}\omega, \tilde{u}_0(\theta_{-t}\omega))|^2 dx + \alpha \int_{\mathbb{R}^n} \rho\left(\frac{|x|^2}{k^2}\right) |\tilde{v}(t, \theta_{-t}\omega, \tilde{v}_0(\theta_{-t}\omega))|^2 dx \leq c\epsilon,$$

where c is independent of ϵ , and hence, for all $t \geq T_5$ and $k \geq R_5$,

$$\int_{|x| \geq \sqrt{2}k} (\beta |\tilde{u}(t, \theta_{-t}\omega, \tilde{u}_0(\theta_{-t}\omega))|^2 dx + \alpha |\tilde{v}(t, \theta_{-t}\omega, \tilde{v}_0(\theta_{-t}\omega))|^2) dx \leq c\epsilon.$$

This completes the proof. $\square$

We now derive uniform estimates on the tails of u and v in $L^2(\mathbb{R}^n)$.

Lemma 4.7. *Assume that $g, h \in L^2(\mathbb{R}^n)$ and (3.4)-(3.7) hold. Let $B = \{B(\omega)\}_{\omega \in \Omega} \in \mathcal{D}$. Then for every $\epsilon > 0$ and P -a.e. $\omega \in \Omega$, there exist $T = T_B(\omega, \epsilon) > 0$ and $R = R(\omega, \epsilon) > 0$ such that, for all $t \geq T$,*

$$\int_{|x| \geq R} (|u(t, \theta_{-t}\omega, u_0(\theta_{-t}\omega))|^2 + |v(t, \theta_{-t}\omega, v_0(\theta_{-t}\omega))|^2) dx \leq \epsilon.$$

Proof. Let T and R be the constants in Lemma 4.6. By (4.64) and (3.10) we have, for all $t \geq T$ and $k \geq R$,

$$\begin{aligned} \int_{|x| \geq R} (|z_1(\omega)|^2 + |z_2(\omega)|^2) dx &= \int_{|x| \geq R} (|\phi_1(x)|^2 |y_1(\omega_1)|^2 + |\phi_2(x)|^2 |y_2(\omega_2)|^2) dx \\ &\leq \frac{\epsilon}{r(\omega)} (|y_1(\omega_1)|^2 + |y_2(\omega_2)|^2) \leq \epsilon. \end{aligned} \quad (4.66)$$

Then by (4.66) and Lemma 4.6, we get that, for all $t \geq T$ and $k \geq R$,

$$\begin{aligned} &\int_{|x| \geq R} (|u(t, \theta_{-t}\omega, u_0(\theta_{-t}\omega))|^2 + |v(t, \theta_{-t}\omega, v_0(\theta_{-t}\omega))|^2) dx \\ &= \int_{|x| \geq R} (|\tilde{u}(t, \theta_{-t}\omega, \tilde{u}_0(\theta_{-t}\omega)) + z_1(\omega)|^2 + |\tilde{v}(t, \theta_{-t}\omega, \tilde{v}_0(\theta_{-t}\omega)) + z_2(\omega)|^2) dx \\ &\leq 2 \int_{|x| \geq R} (|\tilde{u}(t, \theta_{-t}\omega, \tilde{u}_0(\theta_{-t}\omega))|^2 + |\tilde{v}(t, \theta_{-t}\omega, \tilde{v}_0(\theta_{-t}\omega))|^2 + |z_1(\omega)|^2 + |z_2(\omega)|^2) dx \leq 4\epsilon, \end{aligned}$$

which completes the proof. $\square$

Note that equation (3.14) has no any smoothing effect on the solutions, that is, if the initial condition $\tilde{v}_0(\omega)$ is in $L^2(\mathbb{R}^n)$ only, then for any $t \geq 0$, $\tilde{v}(t, \omega, \tilde{v}_0(\omega))$ only belongs to $L^2(\mathbb{R}^n)$, but not $H^1(\mathbb{R}^n)$. Therefore, the compactness of Sobolev embeddings cannot be used directly to derive the asymptotic compactness of the solution operator. To overcome the difficulty, we need to decompose the solution operator as in the deterministic case. Let $\tilde{v}_1$ and $\tilde{v}_2$ be the solutions of the following problems, respectively,

$$\begin{cases} \frac{d\tilde{v}_1}{dt} + \delta\tilde{v}_1 = 0, \\ \tilde{v}_1(0) = \tilde{v}_0, \end{cases} \quad (4.67)$$

and

$$\begin{cases} \frac{d\tilde{v}_2}{dt} + \delta\tilde{v}_2 = \beta\tilde{u} + h + \beta z_1(\theta_t\omega), \\ \tilde{v}_2(0) = 0. \end{cases} \quad (4.68)$$

Then we find that $\tilde{v} = \tilde{v}_1 + \tilde{v}_2$. Let $(u_0(\omega), v_0(\omega)) \in B(\omega)$ with $\{B(\omega)\}_{\omega \in \Omega} \in \mathcal{D}$, and $\tilde{v}_0(\omega) = v_0(\omega) - z_2(\omega)$. Then the solution $\tilde{v}_1(t, \omega, \tilde{v}_0(\omega))$ of (4.67) satisfies that

$$\|\tilde{v}_1(t, \omega, \tilde{v}_0(\omega))\|^2 = \|\tilde{v}_1(t, \omega, v_0(\omega) - z_2(\omega))\|^2 = e^{-\delta t} \|v_0(\omega) - z_2(\omega)\|^2,$$

and

$$\|\tilde{v}_1(t, \theta_{-t}\omega, v_0(\theta_{-t}\omega) - z_2(\theta_{-t}\omega))\|^2 = e^{-\delta t} \|v_0(\theta_{-t}\omega) - z_2(\theta_{-t}\omega)\|^2 \rightarrow 0, \quad \text{as } t \rightarrow \infty. \quad (4.69)$$

On the other hand, for the solutions of problem (4.68), we have the following estimates.

Lemma 4.8. Assume that $g \in L^2(\mathbb{R}^n)$, $h \in H^1(\mathbb{R}^n)$, and (3.4)-(3.7) hold. Let $B = \{B(\omega)\}_{\omega \in \Omega} \in \mathcal{D}$. Then for P -a.e. $\omega \in \Omega$, there exists $T = T_B(\omega) > 0$ such that for all $t \geq T$,

$$\|\nabla \tilde{v}_2(t, \theta_{-t}\omega, 0)\| \leq c(1 + r(\omega)),$$

where c is a positive deterministic constant and $r(\omega)$ is the tempered function in (3.10).

Proof. Taking the inner product of (4.68) with $\Delta \tilde{v}_2$ in $L^2(\mathbb{R}^n)$, we get that

$$\frac{1}{2} \frac{d}{dt} \|\nabla \tilde{v}_2\|^2 + \delta \|\nabla \tilde{v}_2\|^2 = -\beta(\tilde{u}, \Delta \tilde{v}_2) - (h, \Delta \tilde{v}_2) - \beta(z_1(\theta_t \omega), \Delta \tilde{v}_2). \quad (4.70)$$

Note that the first term on the right-hand side of (4.70) is bounded by

$$\beta(\tilde{u}, \Delta \tilde{v}_2) \leq \beta \|\nabla \tilde{u}\| \|\nabla \tilde{v}_2\| \leq \frac{1}{8} \delta \|\nabla \tilde{v}_2\|^2 + \frac{2\beta^2}{\delta} \|\nabla \tilde{u}\|^2. \quad (4.71)$$

For the second term on the right-hand side of (4.70), we have

$$|(h, \Delta \tilde{v}_2)| \leq \|\nabla h\| \|\nabla \tilde{v}_2\| \leq \frac{1}{8} \delta \|\nabla \tilde{v}_2\|^2 + \frac{2}{\delta} \|h\|_{H^1}^2. \quad (4.72)$$

The last term on the right-hand side of (4.70) is bounded by

$$\beta \|\nabla z_1(\theta_t \omega)\| \|\nabla \tilde{v}_2\| \leq \frac{1}{8} \delta \|\nabla \tilde{v}_2\|^2 + \frac{2\beta^2}{\delta} \|\nabla z_1(\theta_t \omega)\|^2 \leq \frac{1}{8} \delta \|\nabla \tilde{v}_2\|^2 + c|y_1(\theta_t \omega_1)|^2. \quad (4.73)$$

Then it follows from (4.70)-(4.73) that

$$\frac{d}{dt} \|\nabla \tilde{v}_2\|^2 + \delta \|\nabla \tilde{v}_2\|^2 \leq c + c \|\nabla \tilde{u}\|^2 + c|y_1(\theta_t \omega_1)|^2,$$

which implies that, for all $t \geq 0$,

$$\frac{d}{dt} \|\nabla \tilde{v}_2\|^2 + \eta \|\nabla \tilde{v}_2\|^2 \leq c + c \|\nabla \tilde{u}\|^2 + c|y_1(\theta_t \omega_1)|^2. \quad (4.74)$$

Integrating (4.74) on $(0, t)$, we obtain that, for all $t \geq 0$,

$$\begin{aligned} \|\nabla \tilde{v}_2(t, \omega, 0)\|^2 &\leq c \int_0^t e^{\eta(s-t)} ds + c \int_0^t e^{\eta(s-t)} \|\nabla \tilde{u}(s, \omega, \tilde{u}_0(\omega))\|^2 ds + c \int_0^t e^{\eta(s-t)} |y_1(\theta_s \omega_1)|^2 ds \\ &\leq \frac{c}{\eta} + c \int_0^t e^{\eta(s-t)} \|\nabla \tilde{u}(s, \omega, \tilde{u}_0(\omega))\|^2 ds + c \int_0^t e^{\eta(s-t)} |y_1(\theta_s \omega_1)|^2 ds. \end{aligned} \quad (4.75)$$

Replacing ω by $\theta_{-t}\omega$ in (4.75), by (3.12) and Lemma 4.2 we get that, for all $t \geq 0$,

$$\begin{aligned}
\|\nabla \tilde{v}_2(t, \theta_{-t}\omega, 0)\|^2 &\leq \frac{c}{\eta} + c \int_0^t e^{\eta(s-t)} \|\nabla \tilde{u}(s, \theta_{-t}\omega, \tilde{u}_0(\theta_{-t}\omega))\|^2 ds + c \int_0^t e^{\eta(s-t)} |y_1(\theta_{s-t}\omega_1)|^2 ds \\
&\leq ce^{-\eta t} (\|\tilde{u}_0(\theta_{-t}\omega)\|^2 + \|\tilde{v}_0(\theta_{-t}\omega)\|^2) + c(1 + r(\omega)) + c \int_{-t}^0 e^{\eta\tau} |y_1(\theta_\tau\omega_1)|^2 d\tau. \\
&\leq ce^{-\eta t} (\|\tilde{u}_0(\theta_{-t}\omega)\|^2 + \|\tilde{v}_0(\theta_{-t}\omega)\|^2) + c(1 + r(\omega)) + cr(\omega) \int_{-t}^0 e^{\frac{1}{2}\eta\tau} d\tau. \\
&\leq ce^{-\eta t} (\|\tilde{u}_0(\theta_{-t}\omega)\|^2 + \|\tilde{v}_0(\theta_{-t}\omega)\|^2) + c(1 + r(\omega)),
\end{aligned}$$

which shows that there is $T_1 > 0$ such that for all $t \geq T_1$,

$$\|\nabla \tilde{v}_2(t, \theta_{-t}\omega, 0)\|^2 \leq 2(1 + r(\omega)).$$

The proof is complete. □

5 Random attractors

In this section, we prove the existence of a $\mathcal{D}$ -random attractor for the random dynamical system Φ associated with the stochastic FitzHugh-Nagumo system on R^n . To this end, we first establish the $\mathcal{D}$ -pullback asymptotic compactness of Φ .

In the sequel, for every $t \in \mathbb{R}^+$, $\omega \in \Omega$ and $(u_0, v_0) \in L^2(\mathbb{R}^n) \times L^2(\mathbb{R}^n)$, we denote by

$$\Phi_1(t, \omega, (u_0, v_0)) = (0, \tilde{v}_1(t, \omega, v_0 - z_2(\omega))) \in L^2(\mathbb{R}^n) \times L^2(\mathbb{R}^n),$$

and

$$\Phi_2(t, \omega, (u_0, v_0)) = (u(t, \omega, u_0), \tilde{v}_2(t, \omega, 0) + z_2(\theta_t\omega)) \in L^2(\mathbb{R}^n) \times L^2(\mathbb{R}^n),$$

where $u(t, \omega, u_0)$ is given by (3.15), $\tilde{v}_1(t, \omega, v_0 - z_2(\omega))$ is the solution of (4.67) with $\tilde{v}_1(0) = v_0 - z_2(\omega)$, and $\tilde{v}_2(t, \omega, 0)$ is the solution of (4.68) with $\tilde{v}_2(0) = 0$. Then we find that

$$\Phi(t, \omega, (u_0, v_0)) = \Phi_1(t, \omega, (u_0, v_0)) + \Phi_2(t, \omega, (u_0, v_0)).$$

The $\mathcal{D}$ -pullback asymptotic compactness of Φ is given by the following lemma.

Lemma 5.1. Assume that $g \in L^2(\mathbb{R}^n)$, $h \in H^1(\mathbb{R}^n)$ and (3.4)-(3.7) hold. Then the random dynamical system Φ is $\mathcal{D}$ -pullback asymptotically compact in $L^2(\mathbb{R}^n) \times L^2(\mathbb{R}^n)$; that is, for P -a.e. $\omega \in \Omega$, the sequence $\{\Phi(t_n, \theta_{-t_n}\omega, (u_{0,n}(\theta_{-t_n}\omega), v_{0,n}(\theta_{-t_n}\omega)))\}$ has a convergent subsequence in $L^2(\mathbb{R}^n) \times L^2(\mathbb{R}^n)$ provided $t_n \rightarrow \infty$, $B = \{B(\omega)\}_{\omega \in \Omega} \in \mathcal{D}$ and $(u_{0,n}(\theta_{-t_n}\omega), v_{0,n}(\theta_{-t_n}\omega)) \in B(\theta_{-t_n}\omega)$.

Proof. Suppose that $t_n \rightarrow \infty$, $B = \{B(\omega)\}_{\omega \in \Omega} \in \mathcal{D}$ and $(u_{0,n}(\theta_{-t_n}\omega), v_{0,n}(\theta_{-t_n}\omega)) \in B(\theta_{-t_n}\omega)$. Then by Lemma 4.1 and (4.69), we find that, for P -a.e. $\omega \in \Omega$,

$$\{\Phi_2(t_n, \theta_{-t_n}\omega, (u_{0,n}(\theta_{-t_n}\omega), v_{0,n}(\theta_{-t_n}\omega)))\}_{n=1}^\infty \text{ is bounded in } L^2(\mathbb{R}^n) \times L^2(\mathbb{R}^n).$$

Hence, there is $(\xi_1, \xi_2) \in L^2(\mathbb{R}^n) \times L^2(\mathbb{R}^n)$ such that, up to a subsequence,

$$\Phi_2(t_n, \theta_{-t_n}\omega, (u_{0,n}(\theta_{-t_n}\omega), v_{0,n}(\theta_{-t_n}\omega))) \rightarrow (\xi_1, \xi_2) \text{ weakly in } L^2(\mathbb{R}^n) \times L^2(\mathbb{R}^n). \quad (5.1)$$

In what follows, we prove the weak convergence of (5.1) is actually strong convergence. Given $\epsilon > 0$ by (4.69) and Lemmas 4.6-4.7, we find that there is $T_1 = T_1(B, \omega, \epsilon)$ and $R_1 = R_1(\omega, \epsilon)$ such that for all $t \geq T_1$,

$$\int_{|x| \geq R_1} |\Phi_2(t, \theta_{-t}\omega, (u_0(\theta_{-t}\omega), v_0(\theta_{-t}\omega)))|^2 dx \leq \epsilon. \quad (5.2)$$

Since $t_n \rightarrow \infty$, there is $N_1 = N_1(B, \omega, \epsilon)$ such that $t_n \geq T_1$ for every $n \geq N_1$. Hence it follows from (5.2) that for all $n \geq N_1$,

$$\int_{|x| \geq R_1} |\Phi_2(t_n, \theta_{-t_n}\omega, (u_{0,n}(\theta_{-t_n}\omega), v_{0,n}(\theta_{-t_n}\omega)))|^2 dx \leq \epsilon. \quad (5.3)$$

On the other hand, By Lemmas 4.1, 4.5 and 4.8, there is $T_2 = T_2(B, \omega)$ such that for all $t \geq T_2$,

$$\|\Phi_2(t, \theta_{-t}\omega, (u_0(\theta_{-t}\omega), v_0(\theta_{-t}\omega)))\|_{H^1(\mathbb{R}^n) \times H^1(\mathbb{R}^n)}^2 \leq c(1 + r(\omega)). \quad (5.4)$$

Let $N_2 = N_2(B, \omega)$ be large enough such that $t_n \geq T_2$ for $n \geq N_2$. Then by (5.4) we find that, for all $n \geq N_2$,

$$\|\Phi_2(t_n, \theta_{-t_n}\omega, (u_{0,n}(\theta_{-t_n}\omega), v_{0,n}(\theta_{-t_n}\omega)))\|_{H^1(\mathbb{R}^n) \times H^1(\mathbb{R}^n)}^2 \leq c(1 + r(\omega)). \quad (5.5)$$

Denote by $Q_{R_1} = \{x \in \mathbb{R}^n : |x| \leq R_1\}$. By the compactness of embedding $H^1(Q_{R_1}) \times H^1(Q_{R_1}) \hookrightarrow L^2(Q_{R_1}) \times L^2(Q_{R_1})$, it follows from (5.5) that, up to a subsequence,

$$\Phi_2(t_n, \theta_{-t_n}\omega, (u_{0,n}(\theta_{-t_n}\omega), v_{0,n}(\theta_{-t_n}\omega))) \rightarrow (\xi_1, \xi_2) \text{ strongly in } L^2(Q_{R_1}) \times L^2(Q_{R_1}),$$

which shows that for the given $\epsilon > 0$, there exists $N_3 = N_3(B, \omega, \epsilon)$ such that for all $n \geq N_3$,

$$\|\Phi_2(t_n, \theta_{-t_n}\omega, (u_{0,n}(\theta_{-t_n}\omega), v_{0,n}(\theta_{-t_n}\omega))) - (\xi_1, \xi_2)\|_{L^2(Q_{R_1}) \times L^2(Q_{R_1})}^2 \leq \epsilon. \quad (5.6)$$

Note that $(\xi_1, \xi_2) \in L^2(\mathbb{R}^n) \times L^2(\mathbb{R}^n)$. Therefore there exists $R_2 = R_2(\epsilon)$ such that

$$\int_{|x| \geq R_2} (|\xi_1(x)|^2 + |\xi_2(x)|^2) dx \leq \epsilon. \quad (5.7)$$

Let $R_3 = \max\{R_1, R_2\}$ and $N_4 = \max\{N_1, N_3\}$. We find that

$$\begin{aligned} & \|\Phi_2(t_n, \theta_{-t_n}\omega, (u_{0,n}(\theta_{-t_n}\omega), v_{0,n}(\theta_{-t_n}\omega))) - (\xi_1, \xi_2)\|_{L^2(\mathbb{R}^n) \times L^2(\mathbb{R}^n)}^2 \\ & \leq \int_{|x| \leq R_3} |\Phi_2(t_n, \theta_{-t_n}\omega, (u_{0,n}(\theta_{-t_n}\omega), v_{0,n}(\theta_{-t_n}\omega))) - (\xi_1, \xi_2)|^2 dx \\ & \quad + \int_{|x| \geq R_3} |\Phi_2(t_n, \theta_{-t_n}\omega, (u_{0,n}(\theta_{-t_n}\omega), v_{0,n}(\theta_{-t_n}\omega))) - (\xi_1, \xi_2)|^2 dx \\ & \leq \int_{|x| \leq R_3} |\Phi_2(t_n, \theta_{-t_n}\omega, (u_{0,n}(\theta_{-t_n}\omega), v_{0,n}(\theta_{-t_n}\omega))) - (\xi_1, \xi_2)|^2 dx \\ & \quad + 2 \int_{|x| \geq R_3} |\Phi_2(t_n, \theta_{-t_n}\omega, (u_{0,n}(\theta_{-t_n}\omega), v_{0,n}(\theta_{-t_n}\omega)))|^2 dx + 2 \int_{|x| \geq R_3} (|\xi_1(x)|^2 + |\xi_2(x)|^2) dx \end{aligned}$$

By (5.3), (5.6)-(5.7), it follows from the above that, for all $n \geq N_4$,

$$\|\Phi_2(t_n, \theta_{-t_n}\omega, (u_{0,n}(\theta_{-t_n}\omega), v_{0,n}(\theta_{-t_n}\omega))) - (\xi_1, \xi_2)\|_{L^2(\mathbb{R}^n) \times L^2(\mathbb{R}^n)}^2 \leq 5\epsilon,$$

which shows that

$$\Phi_2(t_n, \theta_{-t_n}\omega, (u_{0,n}(\theta_{-t_n}\omega), v_{0,n}(\theta_{-t_n}\omega))) \rightarrow (\xi_1, \xi_2) \quad \text{strongly in } L^2(\mathbb{R}^n) \times L^2(\mathbb{R}^n). \quad (5.8)$$

On the other hand, we have

$$\begin{aligned} & \|\Phi(t_n, \theta_{-t_n}\omega, (u_{0,n}(\theta_{-t_n}\omega), v_{0,n}(\theta_{-t_n}\omega))) - (\xi_1, \xi_2)\|_{L^2(\mathbb{R}^n) \times L^2(\mathbb{R}^n)} \\ & \leq \|\Phi_2(t_n, \theta_{-t_n}\omega, (u_{0,n}(\theta_{-t_n}\omega), v_{0,n}(\theta_{-t_n}\omega))) - (\xi_1, \xi_2)\|_{L^2(\mathbb{R}^n) \times L^2(\mathbb{R}^n)} \\ & \quad + \|\Phi_1(t_n, \theta_{-t_n}\omega, (u_{0,n}(\theta_{-t_n}\omega), v_{0,n}(\theta_{-t_n}\omega)))\|_{L^2(\mathbb{R}^n) \times L^2(\mathbb{R}^n)} \\ & \leq \|\Phi_2(t_n, \theta_{-t_n}\omega, (u_{0,n}(\theta_{-t_n}\omega), v_{0,n}(\theta_{-t_n}\omega))) - (\xi_1, \xi_2)\|_{L^2(\mathbb{R}^n) \times L^2(\mathbb{R}^n)} \\ & \quad + \|\tilde{v}_1(t_n, \theta_{-t_n}\omega, v_{0,n}(\theta_{-t_n}\omega) - z_2(\theta_{-t_n}\omega))\|_{L^2(\mathbb{R}^n) \times L^2(\mathbb{R}^n)} \end{aligned}$$

$$\begin{aligned}
&\leq \|\Phi_2(t_n, \theta_{-t_n}\omega, (u_{0,n}(\theta_{-t_n}\omega), v_{0,n}(\theta_{-t_n}\omega))) - (\xi_1, \xi_2)\|_{L^2(\mathbb{R}^n) \times L^2(\mathbb{R}^n)} \\
&\quad + e^{-\delta t_n} \|v_{0,n}(\theta_{-t_n}\omega) - z_2(\theta_{-t_n}\omega)\|_{L^2(\mathbb{R}^n)}
\end{aligned} \tag{5.9}$$

Then, by (4.69), (5.8) and (5.9) we find that

$$\|\Phi(t_n, \theta_{-t_n}\omega, (u_{0,n}(\theta_{-t_n}\omega), v_{0,n}(\theta_{-t_n}\omega))) - (\xi_1, \xi_2)\|_{L^2(\mathbb{R}^n) \times L^2(\mathbb{R}^n)} \rightarrow 0,$$

which completes the proof. $\square$

We are now in a position to present our main result: the existence of a $\mathcal{D}$ -random attractor for Φ in $L^2(\mathbb{R}^n) \times L^2(\mathbb{R}^n)$.

Theorem 5.2. *Assume that $g \in L^2(\mathbb{R}^n)$, $h \in H^1(\mathbb{R}^n)$ and (3.4)-(3.7) hold. Then the random dynamical system Φ has a unique $\mathcal{D}$ -random attractor in $L^2(\mathbb{R}^n) \times L^2(\mathbb{R}^n)$.*

Proof. Notice that Φ has a closed absorbing set $\{K(\omega)\}_{\omega \in \Omega}$ in $\mathcal{D}$ by Lemma 4.1, and is $\mathcal{D}$ -pullback asymptotically compact in $L^2(\mathbb{R}^n) \times L^2(\mathbb{R}^n)$ by Lemma 5.1. Hence the existence of a unique $\mathcal{D}$ -random attractor for Φ follows from Proposition 2.8 immediately. $\square$

References

- [1] F. Antoci and M. Prizzi, Reaction-Diffusion equations on unbounded thin domains, *Topological Methods in Nonlinear Analysis*, **18** (2001), 283-302.
- [2] F. Antoci and M. Prizzi, Attractors and global averaging of non-autonomous Reaction-Diffusion equations in $\mathbb{R}^n$. *Topological Methods in Nonlinear Analysis*, **20** (2002), 229-259.
- [3] L. Arnold, *Random Dynamical Systems*, Springer-Verlag, 1998.
- [4] J. M. Arrieta, J. W. Cholewa, T. Dlotko, and A. Rodriguez-Bernal, Asymptotic behavior and attractors for Reaction Diffusion equations in unbounded domains, *Nonlinear Analysis*, **56** (2004), 515-554.
- [5] A.V. Babin and M.I. Vishik, *Attractors of Evolution Equations*, North-Holland, Amsterdam, 1992.

- [6] J.M. Ball, Continuity properties and global attractors of generalized semiflows and the Navier-Stokes equations, *J. Nonl. Sci.*, **7** (1997), 475-502.
- [7] J.M. Ball, Global attractors for damped semilinear wave equations, *Discrete Contin. Dyn. Syst.* **10** (2004), 31-52.
- [8] P.W. Bates, H. Lisei and K. Lu, Attractors for stochastic lattice dynamical systems, *Stoch. Dyn.*, **6** (2006), 1-21.
- [9] P.W. Bates, K. Lu and B. Wang, Random Attractors for Stochastic Reaction-Diffusion Equations on Unbounded Domains, *J. Differential Equations* (in press).
- [10] T. Caraballo, J. A. Langa and J. C. Robinson, a stochastic pitchfork bifurcation in a reaction-diffusion equation, *Proc. R. Soc. Lond. A*, **457** (2001), 2041-2061.
- [11] H. Crauel, A. Debussche and F. Flandoli, Random attractors, *J. Dyn. Diff. Eqns.*, **9** (1997), 307-341.
- [12] H. Crauel and F. Flandoli, Attractors for random dynamical systems, *Probab. Th. Re. Fields*, **100** (1994), 365-393.
- [13] F. Flandoli and B. Schmalfuß, Random attractors for the 3D stochastic Navier-Stokes equation with multiplicative noise, *Stoch. Stoch. Rep.*, **59** (1996), 21-45.
- [14] J.M. Ghidaglia, A note on the strong convergence towards attractors for damped forced KdV equations, *J. Differential Equations*, **110** (1994), 356-359.
- [15] O. Goubet and R. Rosa, Asymptotic smoothing and the global attractor of a weakly damped KdV equation on the real line, *J. Differential Equations*, **185** (2002), 25-53.
- [16] J.K. Hale, Asymptotic Behavior of Dissipative Systems, American Mathematical Society, Providence, RI, 1988.
- [17] N. Ju, The H^1 -compact global attractor for the solutions to the Navier-Stokes equations in two-dimensional unbounded domains, *Nonlinearity*, **13** (2000), 1227-1238.

- [18] M. Marion, Finite-dimensional attractors associated with partly dissipative reaction-diffusion systems, *SIAM J. Math. Anal.*, **20** (1989), 816-844.
- [19] F. Morillas and J. Valero, Attractors for Reaction-Diffusion equations in $\mathbb{R}^n$ with continuous nonlinearity, *Asymptotic Analysis*, **44** (2005), 111-130.
- [20] I. Moise and R. Rosa, on the regularity of the global attractor of a weakly damped, forced Korteweg-de Vries equation, *Adv. Differential Equations*, **2** (1997), 257-296.
- [21] I. Moise, R. Rosa and X. Wang, Attractors for non-compact semigroups via energy equations, *Nonlinearity*, **11** (1998), 1369-1393.
- [22] M. Prizzi, Averaging, Conley index continuation and recurrent dynamics in almost-periodic equations, *J. Differential Equations*, **210** (2005), 429-451.
- [23] J.C. Robinson, Infinite-Dimensional Dynamical Systems, Cambridge University Press, Cambridge, UK, 2001.
- [24] A. Rodrigue-Bernal and B. Wang, Attractors for partly dissipative Reaction Diffusion systems in $\mathbb{R}^n$, *J. Math. Anal. Appl.*, **252** (2000), 790-803.
- [25] R. Rosa, The global attractor for the 2D Navier-Stokes flow on some unbounded domains, *Nonlinear Anal.*, **32** (1998), 71-85.
- [26] R. Sell and Y. You, Dynamics of Evolutionary Equations, Springer-Verlag, New York, 2002.
- [27] M. Stanislavova, A. Stefanov and B. Wang, Asymptotic smoothing and attractors for the generalized Benjamin-Bona-Mahony equation on $\mathbb{R}^3$, *J. Differential Equations*, **219** (2005), 451-483.
- [28] C. Sun and C. Zhong, Attractors for the semilinear Reaction-Diffusion equation with distribution derivatives in unbounded domains, *Nonlinear Analysis*, **63** (2005), 49-65.
- [29] R. Temam, Infinite-Dimensional Dynamical Systems in Mechanics and Physics, Springer-Verlag, New York, 1997.

- [30] B. Wang, Attractors for reaction-diffusion equations in unbounded domains, *Physica D*, **128** (1999), 41-52.
- [31] X. Wang, An energy equation for the weakly damped driven nonlinear Schrodinger equations and its applications, *Physica D*, **88** (1995), 167-175.